arXiv:1304.4508v2 [math.AG] 29 May 2013

A classical model for derived critical loci

Dominic Joyce

Abstract

Let $f : U \to \mathbb{A}^1$ be a regular function on a smooth scheme U over a field $\mathbb{K}$. Pantev, Toën, Vaquié and Vezzosi [17, 23] define the 'derived critical locus' $\mathbf{Crit}(f)$, an example of a new class of spaces in derived algebraic geometry, which they call '-1-shifted symplectic derived schemes'.

They show that intersections of algebraic Lagrangians in a smooth symplectic $\mathbb{K}$-scheme, and stable moduli schemes of coherent sheaves on a Calabi–Yau 3-fold over $\mathbb{K}$, are also -1-shifted symplectic derived schemes. Thus, their theory may have applications in algebraic symplectic geometry, and in Donaldson–Thomas theory of Calabi–Yau 3-folds.

This paper defines and studies a new class of spaces we call 'algebraic d-critical loci', which should be regarded as classical truncations of the -1-shifted symplectic derived schemes of [17]. They are simpler than their derived analogues. We also give a complex analytic version of the theory.

In the sequels [3–5, 12] we will apply d-critical loci to motivic and categorified Donaldson–Thomas theory, and to intersections of (derived) complex Lagrangians in complex symplectic manifolds. We will show that the important structures one wants to associate to a derived critical locus — virtual cycles, perverse sheaves and mixed Hodge modules of vanishing cycles, and motivic Milnor fibres — can be defined for oriented d-critical loci.

Contents

1 Introduction **2**

2 The main results **6**
2.1 The sheaves $\mathcal{S}_X, \mathcal{S}_X^0$ and d-critical loci 6
2.2 Examples of sheaves $\mathcal{S}_X, \mathcal{S}_X^0$ and d-critical loci 9
2.3 Comparing critical charts (R, U, f, i) 12
2.4 Canonical bundles of d-critical loci 15
2.5 Orientations on d-critical loci 18

3 Proof of Theorem 2.1 **21**
3.1 Construction of the sheaf $\mathcal{S}_X$ 21
3.2 Parts (a)–(c): properties of $\mathcal{S}_X$ 25
3.3 Modifications to the proof for the algebraic case 27

4 Comparing critical charts (R,U,f,i) 29
4.1 Proof of Proposition 2.12 29
4.2 Proof of Theorem 2.13 30
4.3 Proof of Proposition 2.15 31
4.4 Proof of Proposition 2.16 33
4.5 Proof of Proposition 2.17 35

5 Canonical bundles of d-critical loci 36
5.1 Proof of Proposition 2.18 36
5.2 Proof of Proposition 2.20 39
5.3 Proof of Theorem 2.21 41

References 42

1 Introduction

Pantev, Toën, Vaquié and Vezzosi [17,23] defined a new notion of *derived critical locus*. It is set in the context of Toën and Vezzosi's theory of derived algebraic geometry [20–22], and consists of a quasi-smooth derived scheme $\boldsymbol{X}$ equipped with a -1-shifted symplectic structure ω. In fact Pantev et al. [17] define k-shifted symplectic structures on derived stacks for $k \in \mathbb{Z}$, but the case relevant to this paper is $k=-1$, and derived schemes rather than derived stacks.

The following are examples of -1-shifted symplectic derived schemes:

(a) The critical locus $\mathrm{Crit}(f)$ of a regular function $f : U \to \mathbb{A}^1$ on a smooth $\mathbb{K}$-scheme U.

(b) The intersection $L \cap M$ of smooth Lagrangians L, M in an algebraic symplectic manifold (S,ω).

(c) A moduli scheme $\mathcal{M}$ of stable coherent sheaves on a Calabi–Yau 3-fold.

Parts (b),(c) are the beginning of applications of these structures in symplectic geometry and in Donaldson–Thomas theory of Calabi–Yau 3-folds.

This paper will define and study a new class of geometric objects we call *d-critical loci* (X,s). They are much simpler than -1-shifted symplectic derived schemes, and are entirely 'classical', by which we mean they are defined up to isomorphism in an ordinary category using classical algebraic geometry in the style of Hartshorne [9], rather than being defined up to equivalence in an ∞-category using homotopy theory and derived algebraic geometry as in [20–22].

In fact we give two versions of the theory, *complex analytic d-critical loci* (X,s) in which X is a complex analytic space, and *algebraic d-critical loci* (X,s) in which X is a scheme over a field $\mathbb{K}$. In both cases $s \in H^0(\mathcal{S}_X^0)$ is a global section of a certain sheaf $\mathcal{S}_X^0$ on X, satisfying some local conditions. When we can we give results and/or proofs for both complex analytic and algebraic versions simultaneously, or just briefly indicate the differences between the two.

In the algebraic case there are several topologies we could work with — the Zariski topology, the étale topology, and for an embedding $X \hookrightarrow U$ of a $\mathbb{K}$-scheme X into a smooth $\mathbb{K}$-scheme U, it may be natural to consider the formal completion $\hat{U}$ of U along X, and work Zariski or étale locally on $\hat{U}$. Whenever we can, we will use the Zariski topology. One reason is that Theorem 1.4 below, proved in [5], requires the Zariski rather than the étale topology, as distinct motives in $\mathcal{M}^{\mu}_X[\mathbb{L}^{-1/2}]$ can become equal on an étale open cover of X.

To persuade the reader that d-critical loci are a useful idea, we quote four results from the sequels [3–5, 12] to this paper:

Theorem 1.1 (Bussi, Brav and Joyce [4])**.** *Suppose $(\boldsymbol{X},\omega)$ is a -1-shifted symplectic derived scheme over a field $\mathbb{K}$ in the sense of Pantev et al.* [17], *and let $X = t_0(\boldsymbol{X})$ be the associated classical $\mathbb{K}$-scheme of $\boldsymbol{X}$. Then X extends naturally to an algebraic d-critical locus (X,s). The canonical bundle $K_{X,s}$ from* §2.4 *is naturally isomorphic to the determinant line bundle* $\det(\mathbb{L}_{\boldsymbol{X}})|_{X^{\rm red}}$ *of the cotangent complex $\mathbb{L}_{\boldsymbol{X}}$ of $\boldsymbol{X}$. Zariski locally on X one can reconstruct $(\boldsymbol{X},\omega)$ from (X,s) up to equivalence, but this may not be possible globally.*

That is, there is a (non-full) truncation functor from -1-shifted symplectic derived schemes to algebraic d-critical loci. The theorem implies that examples (a)–(c) above have the structure of d-critical loci.

Theorem 1.2 (Joyce [12])**.** *Suppose (S,ω) is a complex symplectic manifold, and L, M are complex Lagrangian submanifolds in S. Then the intersection $X = L\cap M$, as a complex analytic subspace of S, extends naturally to a complex analytic d-critical locus (X,s). The canonical bundle $K_{X,s}$ from* §2.4 *is naturally isomorphic to $K_L|_{X^{\rm red}}\otimes K_M|_{X^{\rm red}}$. The analogue also holds for algebraic symplectic manifolds over a field* $\mathbb{K}$, *and algebraic d-critical loci.*

Thus, we can use complex analytic d-critical loci in the study of (non-algebraic) complex symplectic manifolds. Note that we could not do this using -1-shifted symplectic derived schemes as in [17], without first extending [17, 20–22] to complex analytic geometry.

In §2.4, for a d-critical locus (X,s) we construct a line bundle $K_{X,s}$ on the reduced complex analytic subspace or subscheme $X^{\rm red}$, called the *canonical bundle* of (X,s), and in §2.5 we define an *orientation* on (X,s) to be a choice of square root $K_{X,s}^{1/2}$ of $K_{X,s}$ on $X^{\rm red}$. If $X = \mathrm{Crit}(f)$ for U a complex manifold and $f: U\to\mathbb{C}$ holomorphic then $K_{X,s}\cong K_U^{\otimes^2}|_{X^{\rm red}}$, so there is a natural square root $K_U|_{X^{\rm red}}$ for $K_{X,s}$. Examples in §2.5 show that orientations need not exist, or be unique. Here are two results on oriented d-critical loci:

Theorem 1.3 (Bussi, Brav, Dupont, Joyce and Szendrői [3])**.** *Let (X,s) be a complex analytic d-critical locus with orientation $K_{X,s}^{1/2}$. Then we construct a $\mathbb{Z}$-perverse sheaf $P^\bullet_{X,s}$, a $\mathscr{D}$-module $D_{X,s}$, and a mixed Hodge module $H^\bullet_{X,s}$ on X. If $(X,s), K_{X,s}^{1/2}$ are locally modelled on* $\mathrm{Crit}(f), K_U|_{\mathrm{Crit}(f)^{\rm red}}$ *for U a complex manifold and $f: U\to\mathbb{C}$ holomorphic, then $P^\bullet_{X,s}, D_{X,s}, H^\bullet_{X,s}$ are modelled on the perverse sheaf, $\mathscr{D}$-module and mixed Hodge module of vanishing cycles of U, f.*

Analogues hold for oriented algebraic d-critical loci (X,s), yielding an algebraic $\mathbb{Z}$-perverse sheaf $P^\bullet_{X,s}$, $\mathscr{D}$-module $D_{X,s}$, and mixed Hodge module $H^\bullet_{X,s}$ on X if X is a $\mathbb{C}$-scheme, and a $\mathbb{Z}_l$-perverse sheaf $P^\bullet_{X,s}$ and a $\mathscr{D}$-module $D_{X,s}$ on X if X is a $\mathbb{K}$-scheme and $l \neq \operatorname{char}\mathbb{K}$ a prime.

Theorem 1.4 (Bussi, Joyce and Meinhardt [5])**.** *Let (X,s) be an algebraic d-critical locus over $\mathbb{K}$ with an orientation $K^{1/2}_{X,s}$. Then we construct a motive $MF_{X,s}$ in a certain ring of motives $\mathcal{M}^{\mu}_X[\mathbb{L}^{-1/2}]$ over X. If $(X,s), K^{1/2}_{X,s}$ are Zariski locally modelled on* $\mathrm{Crit}(f), K_U|_{\mathrm{Crit}(f)^{\mathrm{red}}}$ *for U a smooth $\mathbb{K}$-scheme and $f:U\to\mathbb{A}^1$ regular, then $MF_{X,s}$ is locally modelled on $\mathbb{L}^{-\dim U/2}\big([X]-MF^{\mathrm{mot}}_{U,f}\big)$, where $MF^{\mathrm{mot}}_{U,f}$ is the motivic Milnor fibre of f.*

Theorems 1.3 and 1.4 also have applications to extensions of Donaldson–Thomas theory of Calabi–Yau 3-folds, as in [13–15, 19]. Theorem 1.3 is important for *categorification* of Donaldson–Thomas invariants — defining a graded vector space (the hypercohomology $\mathbb{H}^*(P^\bullet_{X,s})$ of the perverse sheaf) whose dimension is the Donaldson–Thomas invariant — and hence for constructing *cohomological Hall algebras*, following Kontsevich and Soibelman [15]. Theorem 1.4 is helpful for defining *motivic* Donaldson–Thomas invariants, as in [14].

We have explained that d-critical loci are classical truncations of -1-shifted symplectic derived schemes in [17]. There is another geometric structure which is a semiclassical truncation of -1-shifted symplectic derived schemes: Behrend's schemes with symmetric obstruction theories [1], which we now define.

Definition 1.5. Let X be a $\mathbb{K}$-scheme. A *perfect obstruction theory* on X in the sense of Behrend and Fantechi [2] is a morphism $\phi:\mathcal{E}^\bullet\to\mathbb{L}_X$ in the derived category $D(\mathrm{qcoh}(X))$, where $\mathbb{L}_X$ is the cotangent complex of X, satisfying:

(i) $\mathcal{E}^\bullet$ is quasi-isomorphic locally on X to a complex $[\mathcal{F}^{-1}\to\mathcal{F}^0]$ of vector bundles in degrees $-1,0$;

(ii) $h^0(\phi):h^0(\mathcal{E}^\bullet)\to h^0(\mathbb{L}_X)$ is an isomorphism; and

(iii) $h^{-1}(\phi):h^{-1}(\mathcal{E}^\bullet)\to h^{-1}(\mathbb{L}_X)$ is surjective.

Following Behrend [1], we call $\phi:\mathcal{E}^\bullet\to\mathbb{L}_X$ a *symmetric obstruction theory* if we also are given an isomorphism $\theta:\mathcal{E}^\bullet\to\mathcal{E}^{\bullet\vee}[1]$ with $\theta^\vee[1]=\theta$.

If U is a smooth $\mathbb{K}$-scheme and $f:U\to\mathbb{A}^1$ a regular function then $X=\mathrm{Crit}(f)$ has a natural symmetric obstruction theory $\phi:\mathcal{E}^\bullet\to\mathbb{L}_X$ with

$$\mathcal{E}^\bullet=\big[TU|_X\xrightarrow{\partial^2 f|_X}T^*U|_X\big]. \tag{1.1}$$

But Pandharipande and Thomas [16] give examples of schemes X with symmetric obstruction theories with X not locally isomorphic to a critical locus. Schemes with symmetric obstruction theories are the basis of Joyce and Song's theory of Donaldson–Thomas invariants of Calabi–Yau 3-folds [13]. If $(\boldsymbol{X},\omega)$ is a -1-shifted symplectic derived scheme in the sense of Pantev et al. [17], then the classical scheme $X=t_0(\boldsymbol{X})$ has a symmetric obstruction theory $\phi:\mathcal{E}^\bullet\to\mathbb{L}_X$ with $\mathcal{E}^\bullet=i^*(\mathbb{L}_{\boldsymbol{X}})$ and $\theta=i^*(\omega_0)$, where $i:X\hookrightarrow\boldsymbol{X}$ is the inclusion.

We illustrate the relations between these structures in Figure 1.1. The two dotted arrows '$\dashrightarrow$' indicate a construction which works locally, but not globally. That is, given an algebraic d-critical locus (X,s), then Zariski locally on X we can construct both a -1-shifted symplectic derived scheme $(\boldsymbol{X},\omega)$, and a symmetric obstruction theory $\phi:\mathcal{E}^\bullet\to\mathbb{L}_X$, θ, uniquely up to equivalence, but we cannot combine these local models to make $(\boldsymbol{X},\omega)$ or $\mathcal{E}^\bullet,\phi,\theta$ globally on X because of difficulties with gluing 'derived' objects on open covers.


−1-shifted symplectic derived schemes [17]
[4]
schemes with symmetric obstruction theories [1]
algebraic d-critical loci
[1, 2]
χ(X,ν_X)
[5]
[3]
virtual cycles
motivic Milnor fibres
perverse sheaves, 𝒟-modules and mixed Hodge modules
[13, 19]
D–T invariants of Calabi–Yau 3-folds [13, 14, 19]
motivic D–T invariants [14]
categorified D–T theory and cohomological Hall algebras [15]
'Fukaya categories' of complex and algebraic symplectic manifolds

Figure 1.1: Relations between different structures, and applications

If X is a proper $\mathbb{K}$-scheme with obstruction theory $\phi:\mathcal{E}^\bullet\to\mathbb{L}_X$ then Behrend and Fantechi define a *virtual cycle* $[X]^{\mathrm{vir}}$ in Chow homology $A_*(X)$. If the obstruction theory is symmetric, and $\mathbb{K}$ algebraically closed of characteristic zero, then Behrend [1] (see also [13, §4]) shows that $[X]^{\mathrm{vir}}\in A_0(X)$, and

$$\textstyle\int_{[X]^{\mathrm{vir}}}1=\chi(X,\nu_X), \tag{1.2}$$

where ν_X is a $\mathbb{Z}$-valued constructible function on X called the *Behrend function*, which depends only on X as a $\mathbb{K}$-scheme. In particular, $[X]^{\mathrm{vir}}$ is independent of the choice of symmetric obstruction theory on X.

If (X,s) is a proper algebraic d-critical locus, we define the *virtual cycle* of X to be $\chi(X,\nu_X)\in\mathbb{Z}$, as in (1.2). Although we will not do it in this paper, one can define a notion of family of d-critical loci over a base Y, and show that the virtual cycles of a proper family of d-critical loci are locally constant on Y.

Example 2.9 below shows that locally, schemes with symmetric obstruction theories can contain strictly less information than algebraic d-critical loci. On the other hand, Example 2.10 shows that schemes with (symmetric) obstruction theories can contain global, nonlocal information (in the form of a class in $\mathrm{Ext}^2(T^*X,T^*X^\vee)$) which is forgotten by algebraic d-critical loci.

The author and his collaborators tried for some time to construct perverse sheaves, and motivic Milnor fibres, from a scheme with symmetric obstruction theory, but failed, and the author now believes this is not possible. So, one moral of Figure 1.1 is that d-critical loci are more useful than schemes with symmetric obstruction theories for various applications.

Conventions. Throughout $\mathbb{K}$ will be an algebraically closed field with char $\mathbb{K} \neq 2$. In this paper all complex analytic spaces and $\mathbb{K}$-schemes X will be assumed to be locally of finite type, as this is necessary for the existence of local embeddings $X \hookrightarrow U$ with U a complex manifold or smooth $\mathbb{K}$-scheme.

Acknowledgements. I would like to thank Oren Ben-Bassat, Dennis Borisov, Chris Brav, Tom Bridgeland, Vittoria Bussi, Stephane Guillermou, Sven Meinhardt, Anatoly Preygel, Pierre Schapira, Edward Segal, Balázs Szendrői, and Bertrand Toën for helpful conversations. This research was supported by EPSRC Programme Grant EP/I033343/1.

2 The main results

This section, the heart of the paper, gives our central definitions, the main results, and some examples. The proofs of results stated in §2.1, §2.3 and §2.4 will be deferred until §3, §4 and §5, respectively.

Some good background references on complex analytic spaces and analytic coherent sheaves upon them are Gunning and Rossi [8] and Grauert and Remmert [7]. A good book on $\mathbb{K}$-schemes and sheaves in algebraic geometry is Hartshorne [9]. The relationship between $\mathbb{C}$-schemes and complex analytic spaces is discussed in Hartshorne [9, App. B] and Serre's famous paper [18].

2.1 The sheaves $\mathcal{S}_X, \mathcal{S}_X^0$ and d-critical loci

The key idea of this section, d-critical loci, is explained in Definition 2.3 below. As a preliminary, we need to associate a sheaf $\mathcal{S}_X$ to each complex analytic space X, such that (very roughly) sections of $\mathcal{S}_X$ parametrize different ways of writing X as $\mathrm{Crit}(f)$ for U a complex manifold and $f : U \to \mathbb{C}$ holomorphic. This we do in the next theorem, which will be proved in §3.

Note our convention from §1 that all complex analytic spaces and $\mathbb{K}$-schemes X in this paper are *locally of finite type*, which is necessary for the existence of embeddings $i : X \hookrightarrow U$ for U a complex manifold or smooth $\mathbb{K}$-scheme.

Theorem 2.1. *Let X be a complex analytic space. Then there exists a sheaf $\mathcal{S}_X$ of $\mathbb{C}$-vector spaces on X, unique up to canonical isomorphism, which is uniquely characterized by the following two properties:*

(i) *Suppose U is a complex manifold, R is an open subset in X, and $i : R \hookrightarrow U$ is an embedding of R as a closed complex analytic subspace of U. Then we have an exact sequence of sheaves of $\mathbb{C}$-vector spaces on R:*

$$\begin{tikzcd} 0 \arrow[r] & I_{R,U} \arrow[r] & i^{-1}(\mathcal{O}_U) \arrow[r, "i^\sharp"] & \mathcal{O}_X|_R \arrow[r] & 0, \end{tikzcd} \tag{2.1}$$

where $\mathcal{O}_X, \mathcal{O}_U$ are the sheaves of holomorphic functions on X, U, and $i^\sharp$ is the morphism of sheaves of $\mathbb{C}$-algebras on R induced by i, which is surjective as i is an embedding, and $I_{R,U} = \mathrm{Ker}(i^\sharp)$ is the sheaf of ideals in $i^{-1}(\mathcal{O}_U)$ of functions on U near $i(R)$ which vanish on $i(R)$.

There is an exact sequence of sheaves of $\mathbb{C}$-vector spaces on R:

$$0 \longrightarrow \mathcal{S}_X|_R \xrightarrow{\iota_{R,U}} \frac{i^{-1}(\mathcal{O}_U)}{I_{R,U}^2} \xrightarrow{\mathrm{d}} \frac{i^{-1}(T^*U)}{I_{R,U}\cdot i^{-1}(T^*U)}, \tag{2.2}$$

where d maps $f + I_{R,U}^2 \mapsto \mathrm{d}f + I_{R,U}\cdot i^{-1}(T^*U)$.

(ii) *Let $R, U, i, \iota_{R,U}$ and $S, V, j, \iota_{S,V}$ be as in* **(i)** *with $R \subseteq S \subseteq X$, and suppose $\Phi : U \to V$ is holomorphic with $\Phi \circ i = j|_R$ as a morphism of complex analytic spaces $R \to V$. Then the following diagram of sheaves on R commutes:*

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{S}_X|_R \arrow[r, "\iota_{S,V}|_R"] \arrow[d, "\mathrm{id}"] & \frac{j^{-1}(\mathcal{O}_V)}{I_{S,V}^2}\Big|_R \arrow[r, "\mathrm{d}"] \arrow[d, "i^{-1}(\Phi^\sharp)"] & \frac{j^{-1}(T^*V)}{I_{S,V}\cdot j^{-1}(T^*V)}\Big|_R \arrow[d, "i^{-1}(\mathrm{d}\Phi)"] \\
0 \arrow[r] & \mathcal{S}_X|_R \arrow[r, "\iota_{R,U}"] & \frac{i^{-1}(\mathcal{O}_U)}{I_{R,U}^2} \arrow[r, "\mathrm{d}"] & \frac{i^{-1}(T^*U)}{I_{R,U}\cdot i^{-1}(T^*U)}.
\end{tikzcd} \tag{2.3}$$

Here $\Phi : U \to V$ induces $\Phi^\sharp : \Phi^{-1}(\mathcal{O}_V) \to \mathcal{O}_U$ on U, so we have

$$i^{-1}(\Phi^\sharp) : j^{-1}(\mathcal{O}_V)|_R = i^{-1}\circ\Phi^{-1}(\mathcal{O}_V) \longrightarrow i^{-1}(\mathcal{O}_U), \tag{2.4}$$

a morphism of sheaves of $\mathbb{C}$-algebras on R. As $\Phi \circ i = j|_R$, equation (2.4) *maps $I_{S,V}|_R \to I_{R,U}$, and so maps $I_{S,V}^2|_R \to I_{R,U}^2$. Thus* (2.4) *induces the morphism in the second column of* (2.3)*. Similarly, $\mathrm{d}\Phi : \Phi^{-1}(T^*V) \to T^*U$ induces the third column of* (2.3)*.*

These sheaves $\mathcal{S}_X$ also satisfy:

(a) *There is a natural decomposition $\mathcal{S}_X = \mathcal{S}_X^0 \oplus \mathbb{C}_X$, where $\mathbb{C}_X$ is the constant sheaf on X with fibre $\mathbb{C}$, and the subsheaf $\mathcal{S}_X^0 \subset \mathcal{S}_X$ is the kernel of the composition*

$$\mathcal{S}_X \xrightarrow{\beta_X} \mathcal{O}_X \xrightarrow{i_X^\sharp} \mathcal{O}_{X^{\mathrm{red}}}, \tag{2.5}$$

with X^{red} the reduced complex analytic subspace of X, and $i_X : X^{\mathrm{red}} \hookrightarrow X$ the inclusion.

(b) *There are natural exact sequences of sheaves of $\mathbb{C}$-vector spaces on X:*

$$0 \longrightarrow h^{-1}(\mathbb{L}_X) \xrightarrow[\alpha_X]{} \mathcal{S}_X \xrightarrow[\beta_X]{} \mathcal{O}_X \xrightarrow[\mathrm{d}]{} T^*X \cong h^0(\mathbb{L}_X), \tag{2.6}$$

$$0 \longrightarrow h^{-1}(\mathbb{L}_X) \xrightarrow{\alpha_X^0} \mathcal{S}_X^0 \xrightarrow{\beta_X^0} \mathcal{O}_X \xrightarrow{\mathrm{d}\oplus i_X^\sharp} T^*X \oplus \mathcal{O}_{X^{\mathrm{red}}}, \tag{2.7}$$

where $\mathbb{L}_X$ is the cotangent complex and T^*X the cotangent sheaf of X.

(c) *The sheaf* $\mathcal{S}_X^0$ *is canonically isomorphic to the cohomology of the complex*

$$I_{R,U}^2 \xrightarrow{\mathrm{d}} I_{R,U}\cdot i^{-1}(T^*U) \xrightarrow{\mathrm{d}} i^{-1}(\Lambda^2 T^*U). \tag{2.8}$$

With the exception of **(c)**, *the analogue of all the above also holds for schemes over a field* $\mathbb{K}$ *in algebraic geometry, taking* X *to be a* $\mathbb{K}$*-scheme with structure sheaf* $\mathcal{O}_X$ *and reduced* $\mathbb{K}$*-subscheme* X^{red}*, and* $\mathcal{S}_X$ *a sheaf of* $\mathbb{K}$*-vector spaces on* X*, and* $R \subseteq X$ *a Zariski open* $\mathbb{K}$*-subscheme, and* U *a smooth* $\mathbb{K}$*-scheme, and replacing* $\mathbb{C}_X$ *by* $\mathbb{K}_X$*. For* **(c)**, *we must replace* U *by the formal completion* $\hat{U}$ *of* U *along* $i(R)$*, so that the analogue of* (2.8) *is*

$$\begin{gathered} I_{R,\hat{U}}^2 \xrightarrow{\mathrm{d}} I_{R,\hat{U}}\cdot i^{-1}(T^*\hat{U}) \xrightarrow{\mathrm{d}} i^{-1}(\Lambda^2 T^*\hat{U}), \quad \textit{where} \\ i^{-1}(\mathcal{O}_{\hat{U}}) = \lim_{n\to\infty} i^{-1}(\mathcal{O}_U)/I_{R,U}^n, \quad I_{R,\hat{U}} = \lim_{n\to\infty} I_{R,U}/I_{R,U}^n \subset i^{-1}(\mathcal{O}_{\hat{U}}). \end{gathered} \tag{2.9}$$

The next example, which is central to our theory, shows the point of $\mathcal{S}_X$.

Example 2.2. Let U be a complex manifold, $f : U \to \mathbb{C}$ be holomorphic, and $X = \mathrm{Crit}(f)$, as a closed complex analytic subspace of U. Write $i : X \hookrightarrow U$ for the inclusion, and $I_{X,U} \subseteq i^{-1}(\mathcal{O}_U)$ for the sheaf of ideals vanishing on $X \subseteq U$. Then $i^{-1}(f) \in H^0\bigl(i^{-1}(\mathcal{O}_U)\bigr)$ with $\mathrm{d}\bigl(i^{-1}(f)\bigr) \in H^0\bigl(I_{X,U}\cdot i^{-1}(T^*U)\bigr) \subseteq H^0\bigl(i^{-1}(T^*U)\bigr)$, so $i^{-1}(f)+I_{X,U}^2 \in H^0\bigl(i^{-1}(\mathcal{O}_U)/I_{X,U}^2\bigr)$ with $\mathrm{d}\bigl(i^{-1}(f)+I_{X,U}^2\bigr) \in H^0\bigl(i^{-1}(T^*U)/I_{X,U}\cdot i^{-1}(T^*U)\bigr)$. Thus by equation (2.2) with $R = X$, we see there is a unique section $s \in H^0(\mathcal{S}_X)$ with $\iota_{X,U}(s) = i^{-1}(f) + I_{X,U}^2$.

Thus, if we can write $X = \mathrm{Crit}(f)$ for $f : U \to \mathbb{C}$ holomorphic, then we obtain a natural section $s \in H^0(\mathcal{S}_X)$. Essentially $s = f + I_{\mathrm{d}f}^2$, where $I_{\mathrm{d}f} \subseteq \mathcal{O}_U$ is the ideal generated by $\mathrm{d}f$. Note that $f|_X = f + I_{\mathrm{d}f}$, so s determines $f|_X$. Basically, s remembers all of the information about f which makes sense intrinsically on X, rather than on the ambient space U.

We can also explain the decomposition $\mathcal{S}_X = \mathcal{S}_X^0 \oplus \mathbb{C}_X$ in this example. We will see in Example 2.6 that if $X = \mathrm{Crit}(f)$ then f need not be locally constant on X, but f is locally constant on the reduced complex analytic space X^{red}. Since locally constant functions on $X^{\mathrm{red}} \subseteq U$ extend uniquely to locally constant functions on U near X^{red}, after shrinking U we can uniquely write $f = f^0 + c$, where $c : U \to \mathbb{C}$ is locally constant and $f^0 : U \to \mathbb{C}$ has $f^0|_{X^{\mathrm{red}}} = 0$. Then f^0, c correspond to the components of s in $H^0(\mathcal{S}_X^0), H^0(\mathbb{C}_X)$, and $X = \mathrm{Crit}(f^0)$.

The analogue also holds in the algebraic case, with U a smooth $\mathbb{K}$-scheme and $f : U \to \mathbb{A}^1$ a regular function.

We can now define d-critical loci:

Definition 2.3. A (*complex analytic*) *d-critical locus* is a pair (X, s), where X is a complex analytic space, and $s \in H^0(\mathcal{S}_X^0)$ for $\mathcal{S}_X^0$ as in Theorem 2.1, satisfying the condition that for each $x \in X$, there exists an open neighbourhood R of x in X, a complex manifold U, a holomorphic function $f : U \to \mathbb{C}$, and an embedding

$i : R \hookrightarrow U$ of R as a closed complex analytic subspace of U, such that $i(R) = \mathrm{Crit}(f)$ as complex analytic subspaces of U, and $\iota_{R,U}(s|_R) = i^{-1}(f) + I_{R,U}^2$.

Similarly, for $\mathbb{K}$-schemes we define an (*algebraic*) *d-critical locus* to be a pair (X, s), where X is a $\mathbb{K}$-scheme, and $s \in H^0(\mathcal{S}_X^0)$ for $\mathcal{S}_X$ as in Theorem 2.1, such that X may be covered by Zariski open sets $R \subseteq X$ with a closed embedding $i : R \hookrightarrow U$ into a smooth $\mathbb{K}$-scheme U and a regular function $f : U \to \mathbb{A}^1 = \mathbb{K}$, such that $i(R) = \mathrm{Crit}(f)$ as $\mathbb{K}$-subschemes of U, and $\iota_{R,U}(s|_R) = i^{-1}(f) + I_{R,U}^2$.

In both cases we call the quadruple (R, U, f, i) a *critical chart* on (X, s).

Remark 2.4. In Definition 2.3, we could instead have defined a d-critical locus (X, s) to have $s \in H^0(\mathcal{S}_X)$ rather than $s \in H^0(\mathcal{S}_X^0)$, but with the rest of the definition the same. The difference is this: as in Example 2.2, if $X = \mathrm{Crit}(f)$ for holomorphic $f : U \to \mathbb{C}$, then $f|_{X^{\mathrm{red}}}$ is locally constant, and we can write $f = f^0 + c$ uniquely near X in U for $f^0 : U \to \mathbb{C}$ holomorphic with $\mathrm{Crit}(f^0) = X = \mathrm{Crit}(f)$, $f^0|_{X^{\mathrm{red}}} = 0$, and $c : U \to \mathbb{C}$ locally constant with $c|_{X^{\mathrm{red}}} = f|_{X^{\mathrm{red}}}$.

Defining d-critical loci using $s \in H^0(\mathcal{S}_X^0)$, as we have done, corresponds to remembering only the function f^0 near X in U, and forgetting the locally constant function $f|_{X^{\mathrm{red}}} : X^{\mathrm{red}} \to \mathbb{C}$. Equivalently, it corresponds to remembering the closed 1-form $\mathrm{d}f = \mathrm{d}f^0$ on U near $X = (\mathrm{d}f)^{-1}(0)$. In the applications the author has in mind [3–5, 12], taking s in $H^0(\mathcal{S}_X^0)$ rather than $H^0(\mathcal{S}_X)$ is more natural, as there is no canonical value for $f|_{X^{\mathrm{red}}}$ other than $f|_{X^{\mathrm{red}}} = 0$. Also (2.8)–(2.9) give an alternative description for $\mathcal{S}_X^0$ rather than $\mathcal{S}_X$.

As in Hartshorne [9, App. B] and Serre [18], there is an analytification functor from algebraic $\mathbb{C}$-schemes X to complex analytic spaces X^{an}, where the points of X^{an} are the $\mathbb{C}$-points of X. It is easy to show that this extends to d-critical loci, and we leave the details to the reader:

Proposition 2.5. *Let (X, s) be an algebraic d-critical locus over the field $\mathbb{K} = \mathbb{C}$. Then the complex analytic space X^{an} associated to the $\mathbb{C}$-scheme X extends naturally to a complex analytic d-critical locus $(X^{\mathrm{an}}, s^{\mathrm{an}})$.*

2.2 Examples of sheaves $\mathcal{S}_X, \mathcal{S}_X^0$ and d-critical loci

Let U be a complex manifold, $f : U \to \mathbb{C}$ be holomorphic, and $X = \mathrm{Crit}(f)$, as a complex analytic space. Then $f|_X : X \to \mathbb{C}$ is holomorphic, and $\mathrm{d}(f|_X) = 0$ in $H^0(T^*X)$. Experience with calculus on manifolds suggests that if $g : X \to \mathbb{C}$ is holomorphic with $\mathrm{d}g = 0$ in $H^0(T^*X)$ then g is locally constant on X. However, this is true only for *reduced* complex analytic spaces or $\mathbb{K}$-schemes X. Here is an example of a non-reduced critical locus with $f|_X$ not locally constant:

Example 2.6. Define $f : \mathbb{C}^2 \to \mathbb{C}$ by $f(x, y) = x^5 + x^2y^2 + y^5$, and let $X = \mathrm{Crit}(f)$, as a complex analytic space. Then $f|_X \in H^0(\mathcal{O}_X)$. We have $\mathrm{d}(f|_X) = 0 \in H^0(T^*X)$, since $X = \mathrm{d}f^{-1}(0)$. Suppose for a contradiction that f is constant on X near $(0, 0)$. Then we may write

$$f(x,y) = A + \tfrac{\partial f}{\partial x}\, B(x,y) + \tfrac{\partial f}{\partial y}\, C(x,y)$$

on $\mathbb{C}^2$ near $(0,0)$, for some holomorphic functions B, C defined near $(0,0)$ in $\mathbb{C}^2$. That is, we have

$$x^5 + x^2y^2 + y^5 = A + (5x^4 + 2xy^2)\sum_{i,j\geqslant 0} B_{i,j}x^iy^j + (2x^2y + 5y^4)\sum_{i,j\geqslant 0} C_{i,j}x^iy^j.$$

Comparing coefficients of x^5, y^5, x^2y^2 give the equations

$$1 = 5B_{1,0}, \quad 1 = 5C_{0,1}, \quad 1 = 2B_{1,0} + 2C_{0,1},$$

which have no solution. Thus in this case, $f|_X$ is not locally constant on X.

If $X = \mathrm{Crit}(f)$ and X^{red} is the reduced complex analytic subspace of X, then $f|_{X^{\mathrm{red}}}$ is always locally constant. This is why we defined $\mathcal{S}_X^0$ using restriction to X^{red} in Theorem 2.1(a). Combining Theorem 2.1(a),(b) we deduce:

Corollary 2.7. *Suppose X is a complex analytic space, and the following sequence of sheaves of $\mathbb{C}$-vector spaces on X is exact:*

$$0 \longrightarrow \mathbb{C}_X \xrightarrow{\mathrm{inc}} \mathcal{O}_X \xrightarrow{\mathrm{d}} T^*X, \tag{2.10}$$

where $\mathrm{inc} : \mathbb{C}_X \hookrightarrow \mathcal{O}_X$ *is the inclusion of the constant functions into the holomorphic functions. Then $\mathcal{S}_X^0 \cong h^{-1}(\mathbb{L}_X)$, so that $\mathcal{S}_X^0$ is a coherent sheaf on X, and $\mathcal{S}_X \cong h^{-1}(\mathbb{L}_X) \oplus \mathbb{C}_X$. The analogue also holds for $\mathbb{K}$-schemes.*

Now (2.10) is exact if for $g : X \to \mathbb{C}$ a locally defined holomorphic function, $\mathrm{d}g = 0$ implies g is locally constant. In Example 2.6 this fails, so in this example (2.10) is not exact, and $\mathcal{S}_X^0 \not\cong h^{-1}(\mathbb{L}_X)$, and $\mathcal{S}_X^0$ is not a coherent sheaf on X.

Next we consider smooth complex analytic spaces and $\mathbb{K}$-schemes:

Example 2.8. Suppose X is a complex manifold, considered as a complex analytic space. Then in Theorem 2.1 we see that $\mathcal{S}_X \cong \mathbb{C}_X$, the constant sheaf, and $\mathcal{S}_X^0 = 0$, the zero sheaf. To see this, take $R = X = U$ and $i = \mathrm{id}_X : X \to X$ in Theorem 2.1. Then $I_{X,X} = 0$ by (2.1), and $\mathcal{S}_X \cong \mathrm{Ker}\bigl(\mathrm{d} : \mathcal{O}_X \to T^*X\bigr) \cong \mathbb{C}_X$ by (2.2). As $\mathcal{S}_X^0 = 0$ there is a unique global section $s = 0 \in H^0(\mathcal{S}_X^0)$, and $(X, 0)$ is a complex analytic d-critical locus, as it is $\mathrm{Crit}(0 : X \to \mathbb{C})$.

Similarly, if X is a smooth $\mathbb{K}$-scheme then $\mathcal{S}_X \cong \mathbb{K}_X$, and $\mathcal{S}_X^0 = 0$, and $(X, 0)$ is an algebraic d-critical locus.

Our next two examples compare algebraic d-critical loci with symmetric obstruction theories on $\mathbb{K}$-schemes, as defined in Definition 1.5.

Example 2.9. Let $\mathbb{K}$ be a field of characteristic zero, and define X to be the $\mathbb{K}$-scheme $X = \mathrm{Spec}\bigl(\mathbb{K}[z]/(z^n)\bigr)$ for $n \geqslant 2$. Then X has an obvious embedding $i : X \hookrightarrow \mathbb{A}^1 = \mathrm{Spec}\bigl(\mathbb{K}[z]\bigr)$ as the subscheme $z^n = 0$ in $\mathbb{A}^1$. It is a non-reduced point. Using this embedding $X \hookrightarrow \mathbb{A}^1$, from Theorem 2.1(i) we find that

$$\begin{aligned} H^0(\mathcal{S}_X) = \bigl\{ &a_0 + a_{n+1}z^{n+1} + \cdots + a_{2n-1}z^{2n-1} + (z^{2n}) : \\ &a_0, a_{n+1}, \ldots, a_{2n-1} \in \mathbb{K} \bigr\} \cong \mathbb{K}^n, \end{aligned}$$

and $H^0(\mathcal{S}_X^0) \subset H^0(\mathcal{S}_X)$ is the subspace with $a_0 = 0$, isomorphic to $\mathbb{K}^{n-1}$.

Now let $0 \in U \subseteq \mathbb{A}^1$ be open, and suppose $f : U \to \mathbb{A}^1$ is regular with $f(0) = 0$ and $\mathrm{Crit}(f) = i(X)$. Write $a_k = \frac{1}{k!}\frac{\partial^k f}{\partial z^k}(0)$ for $k = 0, 1, \ldots$. Then $f(0) = 0$ gives $a_0 = 0$, and $\mathrm{Crit}(f) = X$ is equivalent to $a_1 = \cdots = a_n = 0$ and $a_{n+1} \neq 0$. The section $s \in H^0(\mathcal{S}_X^0)$ corresponding to f is $f + (z^{2n}) = a_{n+1}z^{n+1} + \cdots + a_{2n-1}z^{2n-1} + (z^{2n})$. From this we see that if $s = a_{n+1}z^{n+1} + \cdots + a_{2n-1}z^{2n-1} + (z^{2n}) \in H^0(\mathcal{S}_X^0)$, then (X, s) is an algebraic d-critical locus if and only if $a_{n+1} \neq 0$.

By equation (1.1), the natural symmetric obstruction theory on $X = \mathrm{Crit}(f)$ is determined by $\frac{\partial^2 f}{\partial z^2}\big|_X$, that is, by $(n+1)na_{n+1}z^{n-1} + \cdots$ modulo (z^n). Thus, in this example, the d-critical locus (X, s) associated to $\mathrm{Crit}(f)$ records the first $n-1$ coefficients $a_{n+1}, a_{n+2}, \ldots, a_{2n-1}$ in the power series expansion of $f(z) = a_{n+1}z^{n+1} + a_{n+2}z^{n+2} + \cdots$ at 0, but the symmetric obstruction theory $\phi : \mathcal{E}^\bullet \to \mathbb{L}_X$, θ records only the first coefficient a_{n+1}. Hence (at least in this case), *the algebraic d-critical locus remembers more information, locally, than the symmetric obstruction theory.*

Example 2.10. Let $t : U \to \mathbb{A}^1$ be a smooth proper morphism of $\mathbb{K}$-schemes of relative dimension 2, whose fibres U_t for $t \in \mathbb{A}^1$ are $K3$ surfaces. Set $X = U_0 \subset U$, and regard U, t as a 1-parameter family of deformations of X.

We wish to compare $\mathrm{Crit}(t^2 : U \to \mathbb{A}^1)$ and $\mathrm{Crit}(0 : X \to \mathbb{A}^1)$ in the categories (or higher categories) of:

- (i) classical $\mathbb{K}$-schemes;
- (ii) algebraic d-critical loci;
- (iii) schemes with perfect obstruction theories [2];
- (iv) schemes with symmetric obstruction theories [1]; and
- (v) -1-shifted symplectic derived schemes [17].

We will see that the two are isomorphic in (i),(ii), but not equivalent in (iii)–(v). For (i), $\mathrm{Crit}(t^2 : U \to \mathbb{A}^1)$ and $\mathrm{Crit}(0 : X \to \mathbb{A}^1)$ are both X as classical schemes. For (ii), as X is smooth $\mathcal{S}_X^0 = 0$, so $\mathrm{Crit}(t^2 : U \to \mathbb{A}^1)$ and $\mathrm{Crit}(0 : X \to \mathbb{A}^1)$ are both $(X, 0)$ as algebraic d-critical loci.

For (iii), write $\phi : \mathcal{E}^\bullet \to \mathbb{L}_X$ and $\psi : \mathcal{F}^\bullet \to \mathbb{L}_X$ for the obstruction theories from $\mathrm{Crit}(t^2 : U \to \mathbb{A}^1)$ and $\mathrm{Crit}(0 : X \to \mathbb{A}^1)$. Then (1.1) gives

$$\mathcal{E}^\bullet = \big[TU|_X \xrightarrow{\partial^2(t^2)|_X} T^*U|_X\big], \qquad \mathcal{F}^\bullet = \big[TX \xrightarrow{\;\;0\;\;} T^*X|_X\big].$$

We want to know whether $\mathcal{E}^\bullet \cong \mathcal{F}^\bullet$ in $D(\mathrm{qcoh}(X))$. Now $\tau_{\leqslant -1}(\mathcal{E}^\bullet) \cong TX[1]$ and $\tau_{\geqslant 0}(\mathcal{E}^\bullet) \cong T^*X$, so we have a distinguished triangle in $D(\mathrm{qcoh}(X))$:

$$\begin{tikzcd} \cdots \arrow[r] & T^*X[-1] \arrow[r, "\alpha"] & TX[1] \arrow[r] & \mathcal{E}^\bullet \arrow[r] & T^*X \arrow[r] & \cdots . \end{tikzcd}$$

That is, $\mathcal{E}^\bullet$ is the cone on $\alpha : T^*X[-1] \to TX[1]$ in $D(\mathrm{qcoh}(X))$ for some α in $\mathrm{Ext}^2(T^*X, TX)$. Hence $\mathcal{E}^\bullet \cong \mathcal{F}^\bullet$ if and only if $\alpha = 0$.

The normal bundle ν of X in U, and its dual ν^*, are both isomorphic to $\mathcal{O}_X$ as $t : U \to \mathbb{A}^1$ induces isomorphisms $\nu \cong t^*(T_0\mathbb{A}^1)$, $\nu^* \cong t^*(T_0^*\mathbb{A}^1)$. Hence we have exact sequences

$$\begin{tikzcd}
0 \arrow[r] & TX \arrow[r] & TU|_X \arrow[r] & \mathcal{O}_X \arrow[r] & 0,
\end{tikzcd}$$

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{O}_X \arrow[r] & T^*U|_X \arrow[r] & T^*X \arrow[r] & 0.
\end{tikzcd}$$

Let these correspond to elements $\beta' \in \mathrm{Ext}^1(\mathcal{O}_X, TX)$ and $\beta'' \in \mathrm{Ext}^1(T^*X, \mathcal{O}_X)$. Then $\alpha = \beta' \circ \beta'' \in \mathrm{Ext}^2(T^*X, TX) \cong H^2(TX \otimes TX)$.

Under the isomorphisms $\mathrm{Ext}^1(\mathcal{O}_X, TX) \cong H^1(TX) \cong \mathrm{Ext}^1(T^*X, \mathcal{O}_X)$, we see that β', β'' are both identified with $\beta \in H^1(TX)$, which parametrizes the infinitesimal deformation of $\{U_t : t \in \mathbb{A}^1\}$ at $t = 0$, so that informally $\beta \sim \frac{\mathrm{d}}{\mathrm{d}t}U_t|_{t=0}$. The projection of α from $H^2(TX \otimes TX)$ to $H^2(\Lambda^2 TX) \cong \mathbb{K}$ is β^2.

Let us choose the deformation $t : U \to \mathbb{A}^1$ of $X = U_0$ so that the infinitesimal deformation $\beta \in H^1(TX)$ at $t = 0$ satisfies $\beta^2 \neq 0$ in $H^2(\Lambda^2 TX) \cong \mathbb{K}$, which is possible by well known facts about $K3$ surfaces. Then $\alpha \neq 0$ in $\mathrm{Ext}^2(T^*X, TX)$, and $\mathcal{E}^\bullet \not\cong \mathcal{F}^\bullet$. Hence $\mathrm{Crit}(t^2 : U \to \mathbb{A}^1)$ and $\mathrm{Crit}(0 : X \to \mathbb{A}^1)$ are not equivalent as schemes with perfect obstruction theories, as in (iii), and so a fortiori they are also not equivalent as schemes with symmetric obstruction theories, as in (iv), or as -1-shifted symplectic derived schemes, as in (v).

Observe that $\alpha \in \mathrm{Ext}^2(T^*X, TX)$ which distinguishes the obstruction theories is *global* information, which is *locally trivial*: if $Y \subset X$ is any affine open subset then $\alpha|_Y = 0$ as $\mathrm{Ext}^2(T^*Y, TY) = 0$. Thus (at least in this case), *the (symmetric) obstruction theory remembers global, non-local information which is forgotten by the algebraic d-critical locus.*

This example shows that the dotted arrows '$\dashrightarrow$' in Figure 1.1, which indicate local constructions, cannot be made to work globally.

Using related ideas, the author expects that if (X, s) is an algebraic d-critical locus, then there is an obstruction in $\mathrm{Ext}^3\big(T^*X, (T^*X)^\vee\big)$ to finding a symmetric obstruction theory $\phi : \mathcal{E}^\bullet \to \mathbb{L}_X$, θ on X which is locally modelled on (1.1) when (X, s) is locally modelled on $\mathrm{Crit}(f : U \to \mathbb{C})$. But the author does not know of an example in which this obstruction is nonzero. Finding such an example would show that the truncation functor from -1-shifted symplectic derived schemes to algebraic d-critical loci in [4] is not essentially surjective.

2.3 Comparing critical charts (R, U, f, i)

In §2.1 we defined a d-critical locus (X, s) to admit an open cover by *critical charts* (R, U, f, i), which write (X, s) as $\mathrm{Crit}(f)$ in an open set $R \subset X$. We will treat critical charts like coordinate charts on a manifold. Our analogues of transition functions between coordinate charts are called *embeddings*.

Definition 2.11. Let (X, s) be a d-critical locus (either complex analytic or algebraic), and (R, U, f, i) be a critical chart on (X, s). Let $U' \subseteq U$ be (Zariski) open, and set $R' = i^{-1}(U') \subseteq R$, so that $R' \subseteq R \subseteq X$ are (Zariski) open, and

$i' = i|_{R'} : R' \hookrightarrow U'$, and $f' = f|_{U'} : U' \to \mathbb{C}$ or $\mathbb{A}^1$. Then (R', U', f', i') is also a critical chart on (X, s), and we call it a *subchart* of (R, U, f, i). As a shorthand we write $(R', U', f', i') \subseteq (R, U, f, i)$.

Let (R, U, f, i) and (S, V, g, j) be critical charts on (X, s), with $R \subseteq S \subseteq X$. An *embedding* of (R, U, f, i) in (S, V, g, j) is a locally closed embedding $\Phi : U \hookrightarrow V$ of complex manifolds or $\mathbb{K}$-schemes such that $\Phi \circ i = j|_R : R \to V$ and $f = g \circ \Phi : U \to \mathbb{C}$ or $\mathbb{A}^1$. As a shorthand we write $\Phi : (R, U, f, i) \hookrightarrow (S, V, g, j)$ to mean Φ is an embedding of (R, U, f, i) in (S, V, g, j).

Clearly, if $\Phi : (R, U, f, i) \hookrightarrow (S, V, g, j)$ and $\Psi : (S, V, g, j) \hookrightarrow (T, W, h, k)$ are embeddings, then $\Psi \circ \Phi : (R, U, f, i) \hookrightarrow (T, W, h, k)$ is also an embedding.

If $\Phi : (R, U, f, i) \hookrightarrow (S, V, g, k)$ is an embedding then $\dim U \leqslant \dim V$. Thus, embeddings between critical charts $(R, U, f, i), (S, V, g, j)$ usually go in only one direction, and do not have inverses which are embeddings. The author drew some inspiration for these ideas from the theory of Kuranishi spaces in the work of Fukaya, Oh, Ohta and Ono [6, §A] in symplectic geometry: critical charts (R, U, f, i) are like Kuranishi neighbourhoods on a topological space, and embeddings are like coordinate changes between Kuranishi neighbourhoods.

In the algebraic case, it is sometimes convenient to work with critical charts (S, V, g, j) in which $V \subseteq \mathbb{A}^n$ is Zariski open in an affine space $\mathbb{A}^n$. Every critical chart (R, U, f, i) locally admits embeddings into such a (S, V, g, j). The proof of the next proposition uses the assumption $\operatorname{char} \mathbb{K} \neq 2$ from §1.

Proposition 2.12. *Let (R, U, f, i) be a critical chart on an algebraic d-critical locus (X, s). Then for each $x \in R$ there exists a subchart $(R', U', f', i') \subseteq (R, U, f, i)$ with $x \in R'$ and an embedding $\Phi : (R', U', f', i') \hookrightarrow (S, V, g, j)$ into a critical chart (S, V, g, j) with $V \subseteq \mathbb{A}^n$ Zariski open for some $n \geqslant 0$.*

Given two critical charts $(R, U, f, i), (S, V, g, j)$ on (X, s), there need not exist embeddings between them (or their subcharts) in either direction. So to compare $(R, U, f, i), (S, V, g, j)$, we construct embeddings of subcharts (R', U', f', i'), (S', V', g', j') into a third critical chart (T, W, h, k) with $\dim W \geqslant \dim U, \dim V$.

Theorem 2.13. *Let (X, s) be a d-critical locus (either complex analytic or algebraic), and $(R, U, f, i), (S, V, g, j)$ be critical charts on (X, s). Then for each $x \in R \cap S \subseteq X$ there exist subcharts $(R', U', f', i') \subseteq (R, U, f, i)$, $(S', V', g', j') \subseteq (S, V, g, j)$ with $x \in R' \cap S' \subseteq X$, a critical chart (T, W, h, k) on (X, s), and embeddings $\Phi : (R', U', f', i') \hookrightarrow (T, W, h, k)$, $\Psi : (S', V', g', j') \hookrightarrow (T, W, h, k)$.*

Remark 2.14. To see the point of the definition and theorem, we explain how they will be used. Often, given a d-critical locus (X, s), we want to construct some global object on X by gluing together local data by isomorphisms. Examples include the canonical bundle $K_{X,s}$ in §2.4, perverse sheaves, $\mathscr{D}$-modules and mixed Hodge modules on oriented d-critical loci (X, s) in [3], and motivic Milnor fibres on oriented algebraic d-critical loci (X, s) in [5]. For each of these constructions, we use the following method:

(i) For each critical chart (R, U, f, i) on (X, s), we define a geometric structure $\mathcal{G}_{R,U,f,i}$ on R, with $\mathcal{G}_{R',U',f',i'} = \mathcal{G}_{R,U,f,i}|_{R'}$ for $(R', U', f', i') \subseteq (R, U, f, i)$.

(ii) For each embedding $\Phi : (R,U,f,i) \hookrightarrow (S,V,g,j)$, we define a canonical isomorphism $\Phi_* : \mathcal{G}_{R,U,f,i} \to \mathcal{G}_{S,V,g,j}|_R$. We show Φ_* is independent of Φ, that is, if $\Phi, \tilde{\Phi} : (R,U,f,i) \hookrightarrow (S,V,g,j)$ are embeddings then $\Phi_* = \tilde{\Phi}_*$.

(iii) For embeddings $\Phi : (R,U,f,i) \hookrightarrow (S,V,g,j)$, $\Psi : (S,V,g,j) \hookrightarrow (T,W,h,k)$, we show that $(\Psi \circ \Phi)_* = \Psi_*|_R \circ \Phi_* : \mathcal{G}_{R,U,f,i} \to \mathcal{G}_{T,W,h,k}|_R$.

(iv) Choose critical charts $\bigl\{(R_a,U_a,f_a,i_a) : a \in A\bigr\}$ with $\{R_a : a \in A\}$ an open cover of X. Using (ii),(iii) and Theorem 2.13 we obtain isomorphisms $\iota_{ab} : \mathcal{G}_{R_a,U_a,f_a,i_a}|_{R_a \cap R_b} \to \mathcal{G}_{R_b,U_b,f_b,i_b}|_{R_a \cap R_b}$ for $a,b \in A$, with $\iota_{aa} = \mathrm{id}$ and $\iota_{bc} \circ \iota_{ab}|_{R_a \cap R_b \cap R_c} = \iota_{ac}|_{R_a \cap R_b \cap R_c}$ for $a,b,c \in A$. Thus, provided the geometric structures concerned form a sheaf, there exists $\mathcal{G}$ on X, unique up to canonical isomorphism, with $\mathcal{G}|_{R_a} \cong \mathcal{G}_{R_a,U_a,f_a,i_a}$ for all $a \in A$.

Our next three results say roughly that if $\Phi : (R,U,f,i) \hookrightarrow (S,V,g,j)$ is an embedding of critical charts on (X,s), then locally near $j(R)$ in V we have $V \cong U \times \mathbb{C}^n$ or $V \cong U \times \mathbb{A}^n$ and $g \cong f \boxplus z_1^2 + \cdots + z_n^2$, where $n = \dim V - \dim U$. But in the algebraic case we have to be careful about which topology we mean when we say 'locally'. The complex analytic case is straightforward:

Proposition 2.15. *Let (X,s) be a complex analytic d-critical locus, and $\Phi : (R,U,f,i) \hookrightarrow (S,V,g,j)$ an embedding of critical charts on (X,s). Then for each $x \in R$ there exist open neighbourhoods U', V' of $i(x), j(x)$ in U, V with $\Phi(U') \subseteq V'$, and holomorphic $\alpha : V' \to U$, $\beta : V' \to \mathbb{C}^n$ for $n = \dim V - \dim U$, such that $\alpha \times \beta : V' \to U \times \mathbb{C}^n$ is a biholomorphism with an open subset of $U \times \mathbb{C}^n$, and $\alpha \circ \Phi|_{U'} = \mathrm{id}_{U'}$, $\beta \circ \Phi|_{U'} = 0$, $g|_{V'} = f \circ \alpha + (z_1^2 + \cdots + z_n^2) \circ \beta$.*

For the algebraic case, we give two statements. The first is a direct analogue of Proposition 2.15 for the étale topology, regarding $\imath : U' \to U$, $\jmath : V' \to V$ as étale neighbourhoods of $i(x), j(x)$ in U, V, and $\Phi' : U' \to V'$ with $\Phi \circ \imath = \jmath \circ \Phi'$ as the analogue of $\Phi(U') \subseteq V'$. It will be used in [3].

Recall our convention in §1 that the base field $\mathbb{K}$ of X is algebraically closed with $\mathrm{char}\,\mathbb{K} \neq 2$. Both these assumptions will be used in the proofs of Propositions 2.16 and 2.17, since to define V' we need to take square roots in $\mathbb{K}$, and we need $\mathrm{char}\,\mathbb{K} \neq 2$ to diagonalize quadratic forms over $\mathbb{K}$.

Proposition 2.16. *Let (X,s) be an algebraic d-critical locus, and $\Phi : (R,U,f,i) \hookrightarrow (S,V,g,j)$ an embedding of critical charts on (X,s). Then for each $x \in R$ there exist smooth $\mathbb{K}$-schemes U', V', a point $u' \in U'$, and morphisms $\imath : U' \to U$, $\jmath : V' \to V$, $\Phi' : U' \to V'$, $\alpha : V' \to U$, and $\beta : V' \to \mathbb{A}^n$ for $n = \dim V - \dim U$, such that $\imath(u') = i(x)$, and $\imath : U' \to U$, $\jmath : V' \to V$, $\alpha \times \beta : V' \to U \times \mathbb{A}^n$ are étale, and $\Phi \circ \imath = \jmath \circ \Phi'$, $\alpha \circ \Phi' = \imath$, $\beta \circ \Phi' = 0$, and $g \circ \jmath = f \circ \alpha + (z_1^2 + \cdots + z_n^2) \circ \beta$.*

The second holds with U' a Zariski open neighbourhood of x in U, at the cost of giving a more general form for $g \circ \jmath : V' \to \mathbb{A}^1$. It will be used in [5].

Proposition 2.17. *Let (X,s) be an algebraic d-critical locus, and $\Phi : (R,U,f,i) \hookrightarrow (S,V,g,j)$ an embedding of critical charts on (X,s). Then for each $x \in R$*

there exist a Zariski open neighbourhood U' of $i(x)$ in U, a smooth $\mathbb{K}$-scheme V', and morphisms $\jmath : V' \to V$, $\Phi' : U' \to V'$, $\alpha : V' \to U'$, $\beta : V' \to \mathbb{A}^n$ and $q_1, \ldots, q_n : U' \to \mathbb{A}^1 \setminus \{0\}$ for $n = \dim V - \dim U$, such that $\jmath : V' \to V$ and $\alpha \times \beta : V' \to U' \times \mathbb{A}^n$ are étale, $\Phi|_{U'} = \jmath \circ \Phi'$, $\alpha \circ \Phi' = \mathrm{id}_{U'}$, $\beta \circ \Phi' = 0$, and

$$g \circ \jmath = f \circ \alpha + (q_1 \circ \alpha) \cdot (z_1^2 \circ \beta) + \cdots + (q_n \circ \alpha) \cdot (z_n^2 \circ \beta). \tag{2.11}$$

2.4 Canonical bundles of d-critical loci

Propositions 2.15–2.17 locally describe embeddings $\Phi : (R, U, f, i) \hookrightarrow (S, V, g, j)$ of critical charts on (X, s). We can also associate a piece of global data to Φ, a nondegenerate quadratic form q_{UV} on the pullback $i^*(N_{UV})$ of the normal bundle N_{UV} of $\Phi(U)$ in V.

Proposition 2.18. *Let (X, s) be a d-critical locus (either complex analytic or algebraic), and $\Phi : (R, U, f, i) \hookrightarrow (S, V, g, j)$ be an embedding of critical charts on (X, s). Write N_{UV} for the normal bundle of $\Phi(U)$ in V, regarded as a (holomorphic or algebraic) vector bundle on U in the exact sequence*

$$\begin{tikzcd} 0 \arrow[r] & TU \arrow[r, "\mathrm{d}\Phi"] & \Phi^*(TV) \arrow[r, "\Pi_{UV}"] & N_{UV} \arrow[r] & 0, \end{tikzcd} \tag{2.12}$$

so that $i^*(N_{UV})$ is a vector bundle on $R \subseteq X$. Then there exists a unique $q_{UV} \in H^0(S^2 i^*(N_{UV}^*))$ which is a nondegenerate quadratic form on $i^*(N_{UV})$, with the following property in each case:

- **(a)** *If (X, s) is a complex analytic d-critical locus and $x, U', V', n, \alpha, \beta$ are as in Proposition 2.15, writing $\langle \mathrm{d}z_1, \ldots, \mathrm{d}z_n \rangle_{U'}$ for the trivial vector bundle on U' with basis $\mathrm{d}z_1, \ldots, \mathrm{d}z_n$ and $R' = i^{-1}(U') \subseteq R \subseteq X$, there is a natural isomorphism $\hat{\beta} : \langle \mathrm{d}z_1, \ldots, \mathrm{d}z_n \rangle_{U'} \to N_{UV}^*|_{U'}$ such that*

$$\begin{aligned} &\Phi|_{U'}^*(\mathrm{d}\beta^*) = \Pi_{UV}|_{U'}^* \circ \hat{\beta} : \\ &\qquad \Phi|_{U'}^* \circ \beta^*(T_0^*\mathbb{C}^n) = \langle \mathrm{d}z_1, \ldots, \mathrm{d}z_n \rangle_{U'} \longrightarrow \Phi|_{U'}^*(T^*V), \end{aligned} \tag{2.13}$$

$$\text{with} \quad q_{UV}|_{R'} = i|_{R'}^* \big[(S^2\hat{\beta})(\mathrm{d}z_1 \otimes \mathrm{d}z_1 + \cdots + \mathrm{d}z_n \otimes \mathrm{d}z_n) \big]. \tag{2.14}$$

- **(b)** *If (X, s) is an algebraic d-critical locus and $x, U', V', \iota, \jmath, \Phi', \alpha, \beta, n$ are as in Proposition 2.16, then there is an isomorphism $\hat{\beta} : \langle \mathrm{d}z_1, \ldots, \mathrm{d}z_n \rangle_{U'} \to \iota^*(N_{UV}^*)$ making the following diagram of vector bundles on U' commute:*

$$\begin{tikzcd} \iota^*(N_{UV}^*) \arrow[r, "\iota^*(\Pi_{UV}^*)"'] & \iota^* \circ \Phi^*(T^*V) = \Phi'^* \circ \jmath^*(T^*V) \arrow[d, "\Phi'^*(\mathrm{d}\jmath^*)"'] \\ \langle \mathrm{d}z_1, \ldots, \mathrm{d}z_n \rangle_{U'} = \Phi'^* \circ \beta^*(T_0^*\mathbb{C}^n) \arrow[u, dotted, "\hat{\beta}"'] \arrow[r, "\Phi'^*(\mathrm{d}\beta^*)"] & \Phi'^*(T^*V'), \end{tikzcd} \tag{2.15}$$

and if $R' = R \times_{i,U,\iota} U'$ with projections $\rho : R' \to R$, $i' : R' \to U'$, then

$$\rho^*(q_{UV}) = i'^* \big[(S^2\hat{\beta})(\mathrm{d}z_1 \otimes \mathrm{d}z_1 + \cdots + \mathrm{d}z_n \otimes \mathrm{d}z_n) \big]. \tag{2.16}$$

(c) *If (X,s) is an algebraic d-critical locus and $x, U', V', \jmath, \Phi', \alpha, \beta, n, q_a$ are as in Proposition 2.17, then there is an isomorphism $\hat\beta : \langle \mathrm{d}z_1, \ldots, \mathrm{d}z_n\rangle_{U'} \rightarrow \imath^*(N^*_{UV})$ making the following commute:*

$$\begin{tikzcd}
N^*_{UV}|_{U'} \arrow[rr, "\Pi^*_{UV}|_{U'}"'] & & \Phi|^*_{U'}(T^*V) = \Phi'^* \circ \jmath^*(T^*V) \arrow[d, "\Phi'^*(\mathrm{d}\jmath^*)"'] \\
\langle \mathrm{d}z_1, \ldots, \mathrm{d}z_n\rangle_{U'} = \Phi'^* \circ \beta^*(T^*_0\mathbb{C}^n) \arrow[u, dotted, "\hat\beta"] \arrow[rr, "\Phi'^*(\mathrm{d}\beta^*)"] & & \Phi'^*(T^*V'),
\end{tikzcd} \tag{2.17}$$

and if $R' = i^{-1}(U') \subseteq R \subseteq X$, then

$$q_{UV}|_{R'} = i|^*_{R'}\big[q_1 \cdot (S^2\hat\beta)(\mathrm{d}z_1 \otimes \mathrm{d}z_1) + \cdots + q_n \cdot (S^2\hat\beta)(\mathrm{d}z_n \otimes \mathrm{d}z_n)\big]. \tag{2.18}$$

Now suppose $\Psi : (S,V,g,j) \hookrightarrow (T,W,h,k)$ is another embedding of critical charts, so that $\Psi\circ\Phi : (R,U,f,i) \hookrightarrow (T,W,h,k)$ is also an embedding, and define N_{VW}, q_{VW} and N_{UW}, q_{UW} using Ψ and $\Psi\circ\Phi$ as above. Then there are unique morphisms $\gamma_{UVW}, \delta_{UVW}$ which make the following diagram of vector bundles on U commute, with straight lines exact:

$$\begin{tikzcd}
& & & & & & 0 \arrow[d] & \\
& & 0 \arrow[dr] & & & & TU \arrow[dlll, "\mathrm{d}\Phi"'] \arrow[ddll, "\mathrm{d}(\Psi\circ\Phi)"] & 0 \arrow[l] \\
& & & \Phi^*(TV) \arrow[dll, "\Pi_{UV}"'] \arrow[dr, "\Phi^*(\mathrm{d}\Psi)"] & & & & \\
0 \arrow[r] & N_{UV} \arrow[dl] \arrow[drr, dotted, "\gamma_{UVW}"'] & & & (\Psi\circ\Phi)^*(TW) \arrow[dl, "\Pi_{UW}"] \arrow[ddr, "\Phi^*(\Pi_{VW})"] & & & \\
0 & & & N_{UW} \arrow[dl] \arrow[drr, dotted, "\delta_{UVW}"'] & & & & \\
& & 0 & & & \Phi^*(N_{VW}) \arrow[r] \arrow[dr] & 0 & \\
& & & & & & 0. &
\end{tikzcd} \tag{2.19}$$

Pulling back by i^* gives an exact sequence of vector bundles on $R \subseteq X$:

$$\begin{tikzcd}
0 \arrow[r] & i^*(N_{UV}) \arrow[r, "i^*(\gamma_{UVW})"] & i^*(N_{UW}) \arrow[r, "i^*(\delta_{UVW})"] & j^*(N_{VW})|_R \arrow[r] & 0.
\end{tikzcd} \tag{2.20}$$

Then there is a natural isomorphism of vector bundles on R

$$i^*(N_{UW}) \cong i^*(N_{UV}) \oplus j^*(N_{VW})|_R, \tag{2.21}$$

compatible with the exact sequence (2.20)*, which identifies*

$$\begin{gathered} q_{UW} \cong q_{UV} \oplus q_{VW} \oplus 0 \qquad \textit{under the splitting} \\ S^2(i^*(N^*_{UW})) \cong S^2(i^*(N^*_{UV})) \oplus S^2(j^*(N^*_{VW}))|_R \oplus i^*(N^*_{UV}) \otimes j^*(N^*_{VW})|_R. \end{gathered} \tag{2.22}$$

Using N_{UV}, q_{UV} we define an isomorphism of line bundles J_Φ on $R^{\rm red}$:

Definition 2.19. Let $\Phi : (R, U, f, i) \hookrightarrow (S, V, g, j)$ be an embedding of critical charts on a d-critical locus (X, s). Define N_{UV}, q_{UV} as in Proposition 2.18, and set $n = \dim V - \dim U$. Write $R^{\rm red}$ for the reduced complex analytic subspace or reduced $\mathbb{K}$-subscheme of R. Taking top exterior powers in the dual of (2.12) and pulling back to $R^{\rm red}$ using i^* gives an isomorphism of line bundles

$$\rho_{UV} : \bigl(i^*(K_U) \otimes i^*(\Lambda^n N_{UV}^*)\bigr)|_{R^{\rm red}} \xrightarrow{\cong} j^*(K_V)|_{R^{\rm red}}. \tag{2.23}$$

As q_{UV} is a nondegenerate quadratic form on $i^*(N_{UV})$, its determinant $\det(q_{UV})$ is a nonvanishing section of $i^*(\Lambda^n N_{UV}^*)^{\otimes^2}$. Define an isomorphism of line bundles $J_\Phi : i^*\bigl(K_U^{\otimes^2}\bigr)|_{R^{\rm red}} \to j^*\bigl(K_V^{\otimes^2}\bigr)|_{R^{\rm red}}$ on $R^{\rm red}$ by the commutative diagram

$$\begin{tikzcd} i^*\bigl(K_U^{\otimes^2}\bigr)|_{R^{\rm red}} \arrow[r, "\mathrm{id}_{i^*(K_U^2)} \otimes \det(q_{UV})|_{R^{\rm red}}"'] \arrow[dr, bend right=10, "J_\Phi"'] & \bigl(i^*\bigl(K_U^{\otimes^2}\bigr) \otimes i^*\bigl(\Lambda^{\rm top} N_{UV}^*\bigr)^{\otimes^2}\bigr)\big|_{R^{\rm red}} \arrow[d, "\rho_{UV}^{\otimes^2}|_{R^{\rm red}}"'] \\ & j^*\bigl(K_V^{\otimes^2}\bigr)|_{R^{\rm red}}. \end{tikzcd} \tag{2.24}$$

Here are some useful properties of the J_Φ. The proof that J_Φ is independent of Φ needs $R^{\rm red}$ reduced, which is why we restricted to $R^{\rm red}$ in Definition 2.19.

Proposition 2.20. *In Definition* 2.19, *the isomorphism J_Φ is independent of the choice of Φ. That is, if $\Phi, \tilde\Phi : (R, U, f, i) \hookrightarrow (S, V, g, j)$ are embeddings of critical charts then $J_\Phi = J_{\tilde\Phi}$.*

If $\Psi : (S, V, g, j) \hookrightarrow (T, W, h, k)$ is another embedding of critical charts then

$$J_\Psi|_{R^{\rm red}} \circ J_\Phi = J_{\Psi\circ\Phi}. \tag{2.25}$$

We can now define the *canonical bundle* of a d-critical locus.

Theorem 2.21. *Let (X, s) be a d-critical locus (either complex analytic or algebraic), and $X^{\rm red} \subseteq X$ the associated reduced complex analytic space or reduced $\mathbb{K}$-scheme. Then there exists a (holomorphic or algebraic) line bundle $K_{X,s}$ on $X^{\rm red}$ which we call the* ***canonical bundle*** *of (X, s), which is natural up to canonical isomorphism, and is characterized by the following properties:*

- **(i)** *If (R, U, f, i) is a critical chart on (X, s), there is a natural isomorphism*

$$\iota_{R,U,f,i} : K_{X,s}|_{R^{\rm red}} \longrightarrow i^*\bigl(K_U^{\otimes^2}\bigr)|_{R^{\rm red}}, \tag{2.26}$$

  where $K_U = \Lambda^{\dim U} T^*U$ is the canonical bundle of U in the usual sense.

- **(ii)** *Let $\Phi : (R, U, f, i) \hookrightarrow (S, V, g, j)$ be an embedding of critical charts on (X, s), and let J_Φ be as in* (2.24). *Then*

$$\iota_{S,V,g,j}|_{R^{\rm red}} = J_\Phi \circ \iota_{R,U,f,i} : K_{X,s}|_{R^{\rm red}} \longrightarrow j^*\bigl(K_V^{\otimes^2}\bigr)\big|_{R^{\rm red}}. \tag{2.27}$$

Remark 2.22. As in Theorem 1.1 proved in [4], if (X,s) is the truncation of a -1-shifted symplectic derived scheme $(\boldsymbol{X},\omega)$ in the sense of Pantev et al. [17], then $K_{X,s}\cong\det(\mathbb{L}_{\boldsymbol{X}})|_{X^{\rm red}}$. So $K_{X,s}$ is isomorphic to the canonical bundle of the derived scheme $\boldsymbol{X}$ in this case, which is why we call it a canonical bundle.

2.5 Orientations on d-critical loci

The next two definitions will be important in the sequels [3–5,12]. For examples of results on oriented d-critical loci, see Theorems 1.3 and 1.4 above.

Definition 2.23. Let (X,s) be a d-critical locus (either complex analytic or algebraic), and $K_{X,s}$ its canonical bundle from Theorem 2.21. An *orientation* on (X,s) is a choice of square root line bundle $K_{X,s}^{1/2}$ for $K_{X,s}$ on $X^{\rm red}$. That is, an orientation is a (holomorphic or algebraic) line bundle L on $X^{\rm red}$, together with an isomorphism $L^{\otimes^2}=L\otimes L\cong K_{X,s}$. A d-critical locus with an orientation will be called an *oriented d-critical locus*.

We can express orientations in terms of *principal* $\mathbb{Z}_2$*-bundles*.

Definition 2.24. Let (X,s) be a d-critical locus. For each embedding of critical charts $\Phi:(R,U,f,i)\hookrightarrow(S,V,g,j)$ on (X,s), define a principal $\mathbb{Z}_2$-bundle $\pi_\Phi:P_\Phi\rightarrow R$ over R to be the bundle of square roots of the isomorphism J_Φ in (2.24). That is, local sections $s_\alpha:R\rightarrow P_\Phi$ correspond to local isomorphisms $\alpha:i^*(K_U)|_{R^{\rm red}}\rightarrow j^*(K_V)|_{R^{\rm red}}$ with $\alpha\otimes\alpha=J_\Phi$. Note that Proposition 2.20 implies that P_Φ is independent of the choice of Φ, that is, if $\Phi,\tilde\Phi:(R,U,f,i)\hookrightarrow(S,V,g,j)$ are embeddings then $P_\Phi=P_{\tilde\Phi}$.

If $\Psi:(S,V,g,j)\hookrightarrow(T,W,h,k)$ is another embedding of critical charts then (2.25) implies that there is a canonical isomorphism

$$\Xi_{\Psi,\Phi}:P_{\Psi\circ\Phi}\xrightarrow{\cong}P_\Psi|_R\otimes_{\mathbb{Z}_2}P_\Phi, \tag{2.28}$$

such that if local isomorphisms $\alpha:i^*(K_U)|_{R^{\rm red}}\rightarrow j^*(K_V)|_{R^{\rm red}}$, $\beta:j^*(K_V)|_{R^{\rm red}}\rightarrow k^*(K_W)|_{R^{\rm red}}$, $\gamma:i^*(K_U)|_{R^{\rm red}}\rightarrow k^*(K_W)|_{R^{\rm red}}$ with $\alpha\otimes\alpha=J_\Phi$, $\beta\otimes\beta=J_\Psi|_{R^{\rm red}}$, $\gamma\otimes\gamma=J_{\Psi\circ\Phi}$ correspond to local sections $s_\alpha:R\rightarrow P_\Phi$, $s_\beta:R\rightarrow P_\Psi|_R$, $s_\gamma:R\rightarrow P_{\Phi\circ\Phi}$, then $\Xi_{\Psi,\Phi}(s_\gamma)=s_\beta\otimes_{\mathbb{Z}_2}s_\alpha$ if and only if $\gamma=\beta\circ\alpha$, where $\gamma=\beta\circ\alpha$ is possible by (2.25).

Now let $K_{X,s}^{1/2}$ be a choice of orientation on (X,s), as in Definition 2.23. For each critical chart (R,U,f,i) on (X,s), define a principal $\mathbb{Z}_2$-bundle $\pi_{R,U,f,i}:Q_{R,U,f,i}\rightarrow R$ to be the bundle of square roots of the isomorphism $\iota_{R,U,f,i}$ in (2.26). That is, local sections $s_\beta:R\rightarrow Q_{R,U,f,i}$ correspond to local isomorphisms $\beta:K_{X,s}^{1/2}|_{R^{\rm red}}\rightarrow i^*(K_U)|_{R^{\rm red}}$ with $\beta\otimes\beta=\iota_{R,U,f,i}$.

Given an orientation $K_{X,s}^{1/2}$ and an embedding $\Phi:(R,U,f,i)\hookrightarrow(S,V,g,j)$, we have principal $\mathbb{Z}_2$-bundles $\pi_\Phi:P_\Phi\rightarrow R$, $\pi_{R,U,f,i}:Q_{R,U,f,i}\rightarrow R$ and $\pi_{S,V,g,j}:Q_{S,V,g,j}\rightarrow S$. Then there is a natural isomorphism of principal $\mathbb{Z}_2$-bundles

$$\Lambda_\Phi:Q_{S,V,g,j}|_R\xrightarrow{\cong}P_\Phi\otimes_{\mathbb{Z}_2}Q_{R,U,f,i} \tag{2.29}$$

on R, defined as follows: local isomorphisms

$$\alpha : i^*(K_U)|_{R^{\rm red}} \longrightarrow j^*(K_V)|_{R^{\rm red}}, \quad \beta : K_{X,s}^{1/2}|_{R^{\rm red}} \longrightarrow i^*(K_U)|_{R^{\rm red}},$$
$$\text{and} \qquad \gamma : K_{X,s}^{1/2}|_{R^{\rm red}} \longrightarrow j^*(K_V)|_{R^{\rm red}}$$

with $\alpha\otimes\alpha = i|_{R^{\rm red}}^*(J_\Phi)$, $\beta\otimes\beta=\iota_{R,U,f,i}$ and $\gamma\otimes\gamma=\iota_{S,V,g,j}|_{R^{\rm red}}$ correspond to local sections $s_\alpha : R\rightarrow i^*(P_\Phi)$, $s_\beta : R\rightarrow Q_{R,U,f,i}$ and $s_\gamma : R\rightarrow Q_{S,V,g,j}|_R$. Equation (2.27) shows that $\gamma=\alpha\circ\beta$ is a possible solution for γ, and we define Λ_Φ in (2.29) such that $\Lambda_\Phi(s_\gamma)=s_\alpha\otimes_{\mathbb{Z}_2} s_\beta$ if and only if $\gamma=\alpha\circ\beta$. Note that Λ_Φ is independent of the choice of Φ, as J_Φ, P_Φ are.

If $\Psi : (S,V,g,j)\hookrightarrow (T,W,h,k)$ is another embedding of critical charts then it is easy to check that the following diagram commutes:

$$\begin{tikzcd}
Q_{T,W,h,k}|_R \arrow[rr, "\Lambda_{\Psi\circ\Phi}"'] \arrow[d, "\Lambda_\Psi|_R"] & & P_{\Psi\circ\Phi}\otimes_{\mathbb{Z}_2} Q_{R,U,f,i} \arrow[d, "\Xi_{\Psi,\Phi}\otimes\mathrm{id}_{Q_{R,U,f,i}}"'] \\
\bigl(P_\Psi\otimes_{\mathbb{Z}_2} Q_{S,V,g,j}\bigr)\big|_R \arrow[rr, "\mathrm{id}_{P_\Psi|_R}\otimes\Lambda_\Phi"] & & P_\Psi|_R\otimes_{\mathbb{Z}_2} P_\Phi\otimes_{\mathbb{Z}_2} Q_{R,U,f,i}.
\end{tikzcd} \tag{2.30}$$

Proposition 2.25. *Let (X,s) be a d-critical locus. Then Definition* 2.24 *induces an isomorphism between isomorphism classes of orientations $K_{X,s}^{1/2}$ on (X,s), and isomorphism classes of the following collections of data:*

- **(a)** *For each critical chart (R,U,f,i) on (X,s), a choice of principal $\mathbb{Z}_2$-bundle $\pi_{R,U,f,i} : Q_{R,U,f,i}\rightarrow R$ on R, and*
- **(b)** *For each embedding of critical charts $\Phi : (R,U,f,i)\hookrightarrow(S,V,g,j)$, a choice of isomorphism $\Lambda_\Phi : Q_{S,V,g,j}|_R\rightarrow P_\Phi\otimes_{\mathbb{Z}_2} Q_{R,U,f,i}$ as in* (2.29),

such that (2.30) *commutes for all embeddings $\Phi : (R,U,f,i)\hookrightarrow(S,V,g,j)$, $\Psi : (S,V,g,j)\hookrightarrow(T,W,h,k)$, where $P_\Phi, P_\Psi, P_{\Psi\circ\Phi}, \Xi_{\Psi,\Phi}$ are as in the first part of Definition* 2.24.

The proof of Proposition 2.25 is straightforward. Definition 2.24 shows how to go from an orientation $K_{X,s}^{1/2}$ to a collection of data $Q_{R,U,f,i}, \Lambda_\Phi$. For the converse, given a collection of data $Q_{R,U,f,i}, \Lambda_\Phi$, note that each $Q_{R,U,f,i}$ determines a square root $L_{R,U,f,i}$ for $K_{X,s}|_{R^{\rm red}}$ uniquely up to isomorphism for each critical chart (R,U,f,i), and for an embedding $\Phi : (R,U,f,i)\hookrightarrow(S,V,g,j)$ the isomorphism Λ_Φ determines an isomorphism $i_\Phi : L_{R,U,f,i}\rightarrow L_{S,V,g,j}|_{R^{\rm red}}$, and for $\Phi : (R,U,f,i)\hookrightarrow(S,V,g,j)$, $\Psi : (S,V,g,j)\hookrightarrow(T,W,h,k)$, equation (2.30) commuting implies that $i_{\Psi\circ\Phi}=i_\Psi|_{R^{\rm red}}\circ i_\Phi$. By the sheaf property of line bundles, we can then show there exists $K_{X,s}^{1/2}$ unique up to canonical isomorphism, with isomorphisms $K_{X,s}^{1/2}|_{R^{\rm red}}\cong L_{R,U,f,i}$ for all (R,U,f,i), which are compatible with i_Φ for all Φ. We leave the details to the reader.

Remark 2.26. Let $\Phi : (R,U,f,i)\hookrightarrow(S,V,g,j)$ be an embedding of critical charts on a d-critical locus (X,s). Define N_{UV}, q_{UV} as in Proposition 2.18, and $\pi_\Phi : P_\Phi\rightarrow R$ as in Definition 2.24. Then an alternative interpretation of P_Φ is as *the principal $\mathbb{Z}_2$-bundle of orientations of the nondegenerate quadratic form q_{UV} on the vector bundle $i^*(N_{UV})$ over R.*

Thus, Proposition 2.25 shows that an orientation $K_{X,s}^{1/2}$ on (X,s) is equivalent to giving principal $\mathbb{Z}_2$-bundles $Q_{R,U,f,i}\to R$ for each chart (R,U,f,i) on (X,s), such that $Q_{R,U,f,i}$ and $Q_{S,V,g,j}|_R$ differ by the principal $\mathbb{Z}_2$-bundle of orientations of q_{UV} for each embedding $\Phi:(R,U,f,i)\hookrightarrow(S,V,g,j)$. This is why we chose the term *orientation* for $K_{X,s}^{1/2}$. It is closely relation to the notion of *orientation data* in Kontsevich and Soibelman [14, §5].

Here are some examples of canonical bundles and orientations:

Example 2.27. Let U be a complex manifold, $f:U\to\mathbb{C}$ be holomorphic, and (X,s) be the complex analytic d-critical locus from Example 2.2 with $X=\mathrm{Crit}(f)$. Then Theorem 2.21(i) with $(R,U,f,i)=(X,U,f,\mathrm{inc})$ implies that $K_{X,s}\cong K_U^{\otimes^2}|_{X^{\mathrm{red}}}$. Hence $K_{X,s}$ has a natural square root $K_{X,s}^{1/2}=K_U|_{X^{\mathrm{red}}}$, and (X,s) a natural orientation. The analogue holds for algebraic critical loci.

Example 2.28. Let X be a complex manifold, so that $(X,0)$ is a d-critical locus as in Example 2.8. Then Theorem 2.21(i) with $(R,U,f,i)=(X,X,0,\mathrm{id}_X)$ shows that $K_{X,0}\cong K_X^{\otimes^2}$, where K_X is the usual canonical bundle of X. Again, $(X,0)$ has a natural orientation $K_{X,0}^{1/2}=K_X$.

As we call $K_{X,0}$ the canonical bundle of $(X,0)$, one might have expected $K_{X,0}\cong K_X$. The explanation is that as a derived scheme, $\mathrm{Crit}(0:X\to\mathbb{C})$ is not X, but the shifted cotangent bundle $T^*X[1]$, and the degree -1 fibres of the projection $T^*X[1]\to X$ include an extra factor of K_X in $K_{X,0}$.

Example 2.29. Let X be the non-reduced projective $\mathbb{C}$-scheme

$$\bigl\{[x,y,z]\in\mathbb{CP}^2:z^2=2yz=0\bigr\}.$$

The reduced $\mathbb{C}$-subscheme $X^{\mathrm{red}}\subset X\subset\mathbb{CP}^2$ is the $\mathbb{CP}^1$ defined by $z=0$, and X has only one non-reduced point $[1,0,0]$, with $X\setminus\{[1,0,0]\}\cong\mathbb{C}$ smooth. The open neighbourhood $X\setminus\{[0,1,0]\}$ of [1,0,0] in X is isomorphic as a classical $\mathbb{C}$-scheme to $\mathrm{Crit}(yz^2:\mathbb{C}^2\to\mathbb{C})$, where (y,z) are the coordinates on $\mathbb{C}^2$.

Extend X to an algebraic d-critical locus as follows: on $X\setminus\{[0,1,0]\}$, define s as in Example 2.2 using $X\setminus\{[0,1,0]\}\cong\mathrm{Crit}(yz^2:\mathbb{C}^2\to\mathbb{C})$. But $\mathcal{S}_X^0\equiv 0$ on $X\setminus\{[1,0,0]\}$ by Example 2.8 as $X\setminus\{[1,0,0]\}$ is smooth, so s extends uniquely by zero to all of X. Since (X,s) is modelled on $\mathrm{Crit}(yz^2:\mathbb{C}^2\to\mathbb{C})$ on $X\setminus\{[0,1,0]\}$ and on $\mathrm{Crit}(0:\mathbb{C}\to\mathbb{C})$ on $X\setminus\{[1,0,0]\}$, it is an algebraic d-critical locus.

Theorem 2.21 defines a line bundle $K_{X,s}$ on $X^{\mathrm{red}}\cong\mathbb{CP}^1$. Calculation shows that $K_{X,s}\cong\mathcal{O}_{\mathbb{CP}^1}(-5)$. For the smooth algebraic d-critical locus $(\mathbb{CP}^1,0)$ we have $K_{\mathbb{CP}^1,0}\cong K_{\mathbb{CP}^1}^{\otimes^2}\cong\mathcal{O}_{\mathbb{CP}^1}(-4)$ as in Example 2.28, so the effect of the nonreduced point $[1,0,0]$ in X is to modify $K_{X,s}$ from $\mathcal{O}_{\mathbb{CP}^1}(-4)$ to $\mathcal{O}_{\mathbb{CP}^1}(-5)$. Since -5 is odd, $K_{X,s}$ admits no square root. Thus, (X,s) *is an example of a non-orientable algebraic d-critical locus.* We can also consider (X,s) as a complex analytic d-critical locus, where again it is not orientable.

3 Proof of Theorem 2.1

We now prove Theorem 2.1 from §2.1.

3.1 Construction of the sheaf $\mathcal{S}_X$

Let X be a fixed complex analytic space. In this section we construct the sheaf $\mathcal{S}_X$ in Theorem 2.1, satisfying Theorem 2.1(i),(ii). We will use the following notation. Define a *triple* (R,U,i) to be an open subset $R\subseteq X$, a complex manifold U, and an embedding $i:R\hookrightarrow U$ of R as a closed complex analytic subspace of U, as in Theorem 2.1(i). For such a triple (R,U,i), define the sheaf of ideals $I_{R,U}\subseteq i^{-1}(\mathcal{O}_U)$ as in (2.1). We will also write $I'_{R,U}\subseteq\mathcal{O}_U$ for the sheaf of ideals vanishing on the closed complex analytic subspace $i(R)\subseteq U$.

If (R,U,i) is a triple and $U'\subseteq U$ is open, set $R':=i^{-1}(U')\subseteq R$ and $i':=i|_{R'}:R'\hookrightarrow U'$. Then (R',U',i') is another triple, which we call a *subtriple* of (R,U,i), and write as $(R',U',i')\subseteq(R,U,i)$.

For each triple (R,U,i), as in (2.2) define $\mathcal{K}_{R,U,i},\kappa_{R,U}$ by the exact sequence of sheaves of $\mathbb{C}$-vector spaces on R:

$$0\longrightarrow\mathcal{K}_{R,U,i}\xrightarrow{\ \kappa_{R,U}\ }\frac{i^{-1}(\mathcal{O}_U)}{I_{R,U}^2}\xrightarrow{\ \mathrm{d}\ }\frac{i^{-1}(T^*U)}{I_{R,U}\cdot i^{-1}(T^*U)}\,. \tag{3.1}$$

That is, $\kappa_{R,U}:\mathcal{K}_{R,U,i}\to i^{-1}(\mathcal{O}_U)/I_{R,U}^2$ is the kernel of $\mathrm{d}:i^{-1}(\mathcal{O}_U)/I_{R,U}^2\to i^{-1}(T^*U)/I_{R,U}\cdot i^{-1}(T^*U)$. The difference between (2.2) and (3.1) is that (2.2) includes an isomorphism $\mathcal{S}_X|_R\cong\mathcal{K}_{R,U,i}$, but we have not yet defined $\mathcal{S}_X$. If $(R',U',i')\subseteq(R,U,i)$ then $\mathcal{K}_{R',U',i'}=\mathcal{K}_{R,U,i}|_{R'}$ and $\kappa_{R',U'}=\kappa_{R,U}|_{R'}$.

Call $\Phi:(R,U,i)\to(S,V,j)$ a *morphism of triples* if $R\subseteq S\subseteq X$, and $\Phi:U\to V$ is holomorphic with $\Phi\circ i=j|_R:R\to V$. As for (2.3), form the commutative diagram of sheaves of $\mathbb{C}$-vector spaces on R:

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{K}_{S,V,j}|_R \arrow[r, "\kappa_{S,V}|_R"] \arrow[d, dotted, "\Phi^*"] & \dfrac{j^{-1}(\mathcal{O}_V)}{I_{S,V}^2}\Big|_R \arrow[r, "\mathrm{d}"] \arrow[d, "i^{-1}(\Phi^\sharp)"] & \dfrac{j^{-1}(T^*V)}{I_{S,V}\cdot j^{-1}(T^*V)}\Big|_R \arrow[d, "i^{-1}(\mathrm{d}\Phi)"] \\
0 \arrow[r] & \mathcal{K}_{R,U,i} \arrow[r, "\kappa_{R,U}"] & \dfrac{i^{-1}(\mathcal{O}_U)}{I_{R,U}^2} \arrow[r, "\mathrm{d}"] & \dfrac{i^{-1}(T^*U)}{I_{R,U}\cdot i^{-1}(T^*U)}\,.
\end{tikzcd} \tag{3.2}$$

Since the right hand square of (3.2) commutes as in (2.3), exactness of the rows implies there is a unique $\Phi^*:\mathcal{K}_{S,V,j}|_R\to\mathcal{K}_{R,U,i}$ making (3.2) commute.

If $(R',U',i')\subseteq(R,U,i)$ and $(S',V',j')\subseteq(S,V,j)$ with $\Phi(U')\subseteq V'\subseteq V$ then setting $\Phi':=\Phi|_{U'}:U'\to V'$, we have $\Phi'^*=\Phi^*|_{R'}$.

If $\Psi:(S,V,j)\to(T,W,k)$ is another morphism then so is $\Psi\circ\Phi:(R,U,i)\to$

(T,W,k), and by considering the diagram

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{K}_{T,W,k}|_R \arrow[r, "\kappa_{T,W}|_R"] \arrow[d, "\Psi^*|_R"] \arrow[dd, bend right, "(\Psi\circ\Phi)^*"'] & \frac{k^{-1}(\mathcal{O}_W)}{I_{T,W}^2}\Big|_R \arrow[r, "\mathrm{d}"] \arrow[d, "j^{-1}(\Psi^\sharp)|_R"] \arrow[dd, bend right, "i^{-1}((\Psi\circ\Phi)^\sharp)"'] & \frac{k^{-1}(T^*W)}{I_{T,W}\cdot k^{-1}(T^*W)}\Big|_R \arrow[d, "j^{-1}(\mathrm{d}\Psi)|_R"] \arrow[dd, bend right, "i^{-1}(\mathrm{d}(\Psi\circ\Phi))"'] \\
0 \arrow[r] & \mathcal{K}_{S,V,j}|_R \arrow[r, "\kappa_{S,V}|_R"] \arrow[d, "\Phi^*"] & \frac{j^{-1}(\mathcal{O}_V)}{I_{S,V}^2}\Big|_R \arrow[r, "\mathrm{d}"] \arrow[d, "i^{-1}(\Phi^\sharp)"] & \frac{j^{-1}(T^*V)}{I_{S,V}\cdot j^{-1}(T^*V)}\Big|_R \arrow[d, "i^{-1}(\mathrm{d}\Phi)"] \\
0 \arrow[r] & \mathcal{K}_{R,U,i} \arrow[r, "\kappa_{R,U}"] & \frac{i^{-1}(\mathcal{O}_U)}{I_{R,U}^2} \arrow[r, "\mathrm{d}"] & \frac{i^{-1}(T^*U)}{I_{R,U}\cdot i^{-1}(T^*U)},
\end{tikzcd}$$

we see that $(\Psi\circ\Phi)^*=\Phi^*\circ\Psi^*|_R$, that is, the morphisms Φ^* in (3.2) are contravariantly functorial. If $(R,U,i)=(S,V,j)$ and $\Phi=\mathrm{id}_U$ then $\Phi^*=\mathrm{id}$.

We begin with three lemmas. The first is the main point of the proof:

Lemma 3.1. *The morphism Φ^* in* (3.2) *is independent of the choice of Φ. That is, if $\Phi,\tilde\Phi:(R,U,i)\to(S,V,j)$ are morphisms of triples then*

$$\Phi^*=\tilde\Phi^*:\mathcal{K}_{S,V,j}|_R\longrightarrow\mathcal{K}_{R,U,i}. \tag{3.3}$$

Proof. If $x\in R\subseteq S\subseteq X$ and α is a local section of $\mathcal{K}_{S,V,j}|_R$ defined near x in R, then $\alpha=f+(I'_{S,V})^2$ for f a local section of $\mathcal{O}_V$ defined near $j(x)$ in V such that $\mathrm{d}f$ is a section of $I'_{S,V}\cdot T^*V\subseteq T^*V$ near $j(x)$ in V. Then $\Phi^*(\alpha)=f\circ\Phi+(I'_{R,U})^2$ and $\tilde\Phi^*(\alpha)=f\circ\tilde\Phi+(I'_{R,U})^2$ near x in R.

Choose holomorphic coordinates $(z_1,\ldots,z_n)$ on V near $j(x)=\Phi(i(x))=\tilde\Phi(i(x))$. Then by a holomorphic version of Taylor's Theorem we have

$$\begin{aligned}f\circ\tilde\Phi-f\circ\Phi={}&\textstyle\sum_{a=1}^n\bigl(\frac{\partial f}{\partial z_a}\circ\Phi\bigr)\cdot\bigl(z_a\circ\tilde\Phi-z_a\circ\Phi\bigr)\\ &+\textstyle\sum_{a,b=1}^n A_{ab}\bigl(z_a\circ\tilde\Phi-z_a\circ\Phi\bigr)\bigl(z_b\circ\tilde\Phi-z_b\circ\Phi\bigr)\end{aligned} \tag{3.4}$$

near $j(x)$, for some holomorphic $A_{ab}:V\to\mathbb{C}$ defined near $j(x)$.

Since $\Phi\circ i=\tilde\Phi\circ i$ and $I'_{R,U}$ is the ideal in $\mathcal{O}_U$ vanishing on $i(R)$ near $i(x)$, we see that $z_a\circ\tilde\Phi-z_a\circ\Phi\in I'_{R,U}$ on U near $i(x)$ for each $a=1,\ldots,n$. Also $\frac{\partial f}{\partial z_a}\in I'_{S,V}$ near $j(x)$ by choice of α,f, and $g\mapsto g\circ\Phi$ maps $I'_{S,V}\to I'_{R,U}$ near $j(x),i(x)$ as $\Phi\circ i=j|_R$, so $\frac{\partial f}{\partial z_a}\circ\Phi\in I'_{R,U}$ near $i(x)$. Thus each factor $(\cdots)$ on the right hand side of (3.4) lies in $I'_{R,U}$ near $i(x)$, so $f\circ\tilde\Phi-f\circ\Phi\in(I'_{R,U})^2$ near $i(x)$. Therefore

$$\Phi^*(\alpha)=f\circ\Phi+(I'_{R,U})^2=f\circ\tilde\Phi+(I'_{R,U})^2=\tilde\Phi^*(\alpha)$$

near x in R for any local section α of $\mathcal{K}_{S,V,j}|_R$, which proves (3.3). □

Lemma 3.2. *Let* $(R,U,i),(S,V,j)$ *be triples. Then for each* $x\in R\cap S\subseteq X$, *there exists a subtriple* $(R',U',i')\subseteq(R,U,i)$ *with* $x\in R'\subseteq R\cap S$ *and a morphism* $\Phi:(R',U',i')\to(S,V,j)$.

Proof. Choose holomorphic coordinates $(z_1,\dots,z_n)$ on an open neighbourhood $\tilde V$ of $j(x)$ in V, so that $(z_1,\dots,z_n):\tilde V\to\mathbb{C}^n$ is a biholomorphism with an open set $\tilde W\subseteq\mathbb{C}^n$. Let U' be an open neighbourhood of $i(x)$ in U small enough that $R':=i^{-1}(U')\subseteq R\cap j^{-1}(\tilde V)\subseteq R\cap S\subseteq X$. Then $z_a\circ j|_{R'}$ for $a=1,\dots,n$ are morphisms $R'\to\mathbb{C}$. Since $i(R')$ is a closed complex analytic subspace of U' isomorphic to R', and any holomorphic function on $i(R')$ extends locally to U' near $i(x)$, by making R',U' smaller we can suppose there exist holomorphic $f_a:U'\to\mathbb{C}$ with $f_a\circ i|_{R'}=z_a\circ j|_{R'}$ for $a=1,\dots,n$.

Making R',U' smaller again, we can suppose that $(f_1,\dots,f_n):U'\to\mathbb{C}^n$ maps into $\tilde V\subseteq\mathbb{C}^n$. Then there is a unique holomorphic map $\Phi:U'\to\tilde V\subseteq V$ with $z_a\circ\Phi=f_a$ for $a=1,\dots,n$. Hence $z_a\circ\Phi\circ i|_{R'}=f_a\circ i|_{R'}=z_a\circ j|_{R'}$ for $a=1,\dots,n$, which implies that $\Phi\circ i|_{R'}=j|_{R'}$ as $(z_1,\dots,z_n):\tilde V\to\mathbb{C}^n$ is injective. Thus $\Phi:(R',U',i')\to(S,V,j)$ is a morphism of triples. ☐

Lemma 3.3. *Let* $(R,U,i),(S,V,j)$ *be triples. Then there exists a unique isomorphism of sheaves on* $R\cap S$:

$$I_{S,V,j}^{R,U,i}:\mathcal{K}_{S,V,j}|_{R\cap S}\longrightarrow\mathcal{K}_{R,U,i}|_{R\cap S}\tag{3.5}$$

such that if $(R',U',i'),\Phi$ *are as in Lemma 3.2 then* $I_{S,V,j}^{R,U,i}|_{R'}=\Phi^*$. *Also*

$$I_{R,U,i}^{R,U,i}=\mathrm{id}_{\mathcal{K}_{R,U,i}},\qquad I_{R,U,i}^{S,V,j}=\bigl(I_{S,V,j}^{R,U,i}\bigr)^{-1},\tag{3.6}$$

and if (T,W,k) *is another triple then*

$$\begin{gathered}I_{S,V,j}^{R,U,i}|_{R\cap S\cap T}\circ I_{T,W,k}^{S,V,j}|_{R\cap S\cap T}=I_{T,W,k}^{R,U,i}|_{R\cap S\cap T}:\\ \mathcal{K}_{T,W,k}|_{R\cap S\cap T}\longrightarrow\mathcal{K}_{R,U,i}|_{R\cap S\cap T}.\end{gathered}\tag{3.7}$$

Proof. Suppose $x,(R',U',i'),\Phi$ and $\hat x,(\hat R',\hat U',\hat i'),\hat\Phi$ are two possible choices in Lemma 3.2. Then $\Phi:U'\to V$, $\hat\Phi:\hat U'\to V$ induce morphisms

$$\Phi^*:\mathcal{K}_{S,V,j}|_{R'}\longrightarrow\mathcal{K}_{R,U,i}|_{R'}\ \text{ and }\ \hat\Phi^*:\mathcal{K}_{S,V,j}|_{\hat R'}\longrightarrow\mathcal{K}_{R,U,i}|_{\hat R'}.$$

The restrictions $\Phi|_{U'\cap\hat U'}$ and $\hat\Phi|_{U'\cap\hat U'}$ thus induce morphisms

$$\Phi^*|_{R'\cap\hat R'},\hat\Phi^*|_{R'\cap\hat R'}:\mathcal{K}_{S,V,j}|_{R'\cap\hat R'}\longrightarrow\mathcal{K}_{R,U,i}|_{R'\cap\hat R'}.$$

Lemma 3.1 now shows that $\Phi^*|_{R'\cap\hat R'}=\hat\Phi^*|_{R'\cap\hat R'}$.

Thus, Lemma 3.2 shows that for each $x\in R\cap S$ we can choose an open neighbourhood R' of x in $R\cap S$, and a morphism $\Phi^*:\mathcal{K}_{S,V,j}|_{R'}\longrightarrow\mathcal{K}_{R,U,i}|_{R'}$. These open neighbourhoods R' form an open cover of X, and on overlaps $R'\cap\hat R'$ the corresponding morphisms $\Phi^*,\hat\Phi^*$ agree. Therefore by properties of sheaves there is a unique morphism $I^{R,U,i}_{S,V,j}$ in (3.5) such that $I^{R,U,i}_{S,V,j}\big|_{R'} = \Phi^*$ for all $(R',U',i'),\Phi$ as in Lemma 3.2.

To see that $I^{R,U,i}_{R,U,i} = \mathrm{id}_{\mathcal{K}_{R,U,i}}$ as in (3.6), take $(R',U',i') = (R,U,i) = (S,V,j)$ and $\Phi = \mathrm{id}_U$, so that $\Phi^* = \mathrm{id}$.

To prove (3.7), let $(R,U,i),(S,V,j),(T,W,k)$ be triples and $x \in R\cap S\cap T$. Apply Lemma 3.2 twice to get open $i(x)\in U'\subseteq U$ and $j(x)\in V'\subseteq V$ and morphisms $\Phi : (R',U',i') \to (S,V,j)$ and $\Psi : (S',V',j') \to (T,W,k)$. Making U',R' smaller we can suppose that $j(x)\in\Phi(U')\subseteq V'\subseteq V$, so that Φ is also a morphism $(R',U',i')\to(S',V',j')$. Functoriality of the Φ^* now gives $\Phi^*\circ\Psi^*|_{R'} = (\Psi\circ\Phi)^*$. So the defining property of the $I^{R,U,i}_{S,V,j}$ gives

$$I^{R,U,i}_{S,V,j}\big|_{R'} \circ I^{S,V,j}_{T,W,k}\big|_{R'} = I^{R,U,i}_{T,W,k}\big|_{R'}.$$

As we can cover $R\cap S\cap T$ by such open R', equation (3.7) follows.

Finally, applying (3.7) with $(T,W,k) = (R,U,i)$ and using the first equation of (3.6) yields

$$I^{R,U,i}_{S,V,j}\circ I^{S,V,j}_{R,U,i} = I^{R,U,i}_{R,U,i}\big|_{R\cap S} = \mathrm{id}_{\mathcal{K}_{R,U,i}}\big|_{R\cap S}.$$

Exchanging $(R,U,i),(S,V,j)$ proves the second equation of (3.6), and also shows that $I^{R,U,i}_{S,V,j}$ is an isomorphism. The lemma is complete. $\square$

We can now construct the sheaf $\mathcal{S}_X$ in Theorem 2.1. Since X is locally of finite type by our convention in §1, near each $x\in X$ it admits a local embedding $i : X \hookrightarrow U$ into a complex manifold U. Therefore we can choose a family $\big\{(R_a,U_a,i_a) : a\in A\big\}$ of triples such that $\{R_a : a\in A\}$ is an open cover of X. For each $a\in A$ we have a sheaf of $\mathbb{C}$-vector spaces $\mathcal{K}_{R_a,U_a,i_a}$ on R_a from (3.1), and for all $a,b\in A$ we have an isomorphism

$$I^{R_a,U_a,i_a}_{R_b,U_b,i_b} : \mathcal{K}_{R_b,U_b,i_b}|_{R_a\cap R_b} \longrightarrow \mathcal{K}_{R_a,U_a,i_a}|_{R_a\cap R_b}$$

on $R_a\cap R_b$ by (3.5), which satisfy the usual conditions (3.6)–(3.7) on identities, symmetry, and triple overlaps.

Hence by properties of sheaves there exists a sheaf $\mathcal{S}_X$ on X, unique up to canonical isomorphism, with isomorphisms $\iota_{R_a,U_a} : \mathcal{S}_X|_{R_a} \to \mathcal{K}_{R_a,U_a,i_a}$ for each $a\in A$, such that $I^{R_a,U_a,i_a}_{R_b,U_b,i_b}\circ\iota_{R_b,U_b}|_{R_a\cap R_b} = \iota_{R_a,U_a}|_{R_a\cap R_b}$ for all $a,b\in A$. One way to do this is to define an explicit presheaf $\mathcal{PS}_X$ on X using the data $R_a,\mathcal{K}_{R_a,U_a,i_a}, I^{R_a,U_a,i_a}_{R_b,U_b,i_b}$, and take $\mathcal{S}_X$ to be the sheafification of $\mathcal{PS}_X$.

Now suppose (R,U,i) is any triple, as in Theorem 2.1(i). We can construct $\mathcal{S}_X$ using the family $\big\{(R_a,U_a,i_a) : a\in A\big\}\cup\big\{(R,U,i)\big\}$ instead of $\big\{(R_a,U_a,i_a) : a\in A\big\}$, and get the same (not just isomorphic) sheaf $\mathcal{S}_X$ and the same isomorphisms $\iota_{R_a,U_a} : \mathcal{S}_X|_{R_a}\to\mathcal{K}_{R_a,U_a,i_a}$, but now we also have an isomorphism $\iota_{R,U} : \mathcal{S}_X|_R \to \mathcal{K}_{R,U,i}$. Combining this with (3.1) gives the exact sequence (2.2), proving Theorem 2.1(i).

Suppose $\Phi : (R,U,i)\to(S,V,j)$ is any morphism of triples, as in Theorem 2.1(ii), so that $R\subseteq S$. We can construct $\mathcal{S}_X$ using the family $\big\{(R_a,U_a,i_a) : a\in A\big\}\cup\big\{(R,U,i),(S,V,j)\big\}$ instead of $\big\{(R_a,U_a,i_a) : a\in A\big\}$, and get the same sheaf $\mathcal{S}_X$ and isomorphisms ι_{R_a,U_a}, but now we also have isomorphisms

$\iota_{R,U}:\mathcal{S}_X|_R\rightarrow\mathcal{K}_{R,U,i}$ and $\iota_{S,V}:\mathcal{S}_X|_S\rightarrow\mathcal{K}_{S,V,j}$ satisfying $I_{S,V,j}^{R,U,i}\circ\iota_{S,V}|_R=\iota_{R,U}$. Consider the diagram:

$$\begin{tikzcd}
\mathcal{S}_X|_R \arrow[r, "\iota_{S,V}|_R"] \arrow[d, "\mathrm{id}"] & \mathcal{K}_{S,V,j}|_R \arrow[r, "\kappa_{S,V}|_R"] \arrow[d, "I_{S,V,j}^{R,U,i}=\Phi^*"] & \frac{j^{-1}(\mathcal{O}_V)}{I_{S,V}^2}\Big|_R \arrow[r, "\mathrm{d}"] \arrow[d, "i^{-1}(\Phi^\sharp)"] & \frac{j^{-1}(T^*V)}{I_{S,V}\cdot j^{-1}(T^*V)}\Big|_R \arrow[d, "i^{-1}(\mathrm{d}\Phi)"] \\
\mathcal{S}_X|_R \arrow[r, "\iota_{R,U}"] & \mathcal{K}_{R,U,i} \arrow[r, "\kappa_{R,U}"] & \frac{i^{-1}(\mathcal{O}_U)}{I_{R,U}^2} \arrow[r, "\mathrm{d}"] & \frac{i^{-1}(T^*U)}{I_{R,U}\cdot i^{-1}(T^*U)}.
\end{tikzcd} \tag{3.8}$$

Here $I_{S,V,j}^{R,U,i}=\Phi^*$ by Lemma 3.3. The left hand square of (3.8) commutes as $I_{S,V,j}^{R,U,i}\circ\iota_{S,V}|_R=\iota_{R,U}$, and the right hand two squares commute by (3.2). Composing the first two horizontal morphisms in the rows of (3.8) gives (2.3). This proves Theorem 2.1(ii).

3.2 Parts (a)–(c): properties of $\mathcal{S}_X$

Next we prove that the sheaf $\mathcal{S}_X$ constructed in §3.1 satisfies Theorem 2.1(a)–(c). We continue to use the notation of §3.1. For part (a), define $\mathcal{S}_X^0\subset\mathcal{S}_X$ to be the kernel of the morphism $\mathcal{S}_X\rightarrow\mathcal{O}_{X^{\mathrm{red}}}$ in (2.5), where $\beta_X:\mathcal{S}_X\rightarrow\mathcal{O}_X$ is defined using (3.1)–(3.2) and $\mathcal{O}_X|_R\cong i^{-1}(\mathcal{O}_U)/I_{R,U}$.

There is also a natural inclusion $\mathbb{C}_X\hookrightarrow\mathcal{S}_X$, where $\mathbb{C}_X$ is the sheaf of locally constant functions $X\rightarrow\mathbb{C}$: for any triple (R,U,i), the sheaf $\mathbb{C}_U$ of locally constant functions on U is a subsheaf of $\mathcal{O}_U$, so $\bigl(i^{-1}(\mathbb{C}_U)+I_{R,U}^2\bigr)/I_{R,U}^2$ is a subsheaf of $i^{-1}(\mathcal{O}_U)/I_{R,U}^2$ which lies in the kernel of d in (2.2), and so lifts to a subsheaf of $\mathcal{S}_X|_R$, which is isomorphic to $\mathbb{C}_X|_R$, as locally constant functions on X lift locally uniquely to locally constant functions on U near $i(X)$.

It remains to show that $\mathcal{S}_X=\mathcal{S}_X^0\oplus\mathbb{C}_X$. To see this, suppose (R,U,i) is a triple and s is a local section of $\mathcal{S}_X$ near $x\in R$, so that $\iota_{R,U}(s|_R)=f+(I'_{R,U})^2$ for f a local section of $\mathcal{O}_U$ defined near $i(x)$ in U with $\mathrm{d}f$ a local section of $I'_{R,U}\cdot T^*U$. Then $f|_{i(X^{\mathrm{red}})}$ is locally constant on $i(X^{\mathrm{red}})$ near $i(x)$, so $f|_{i(X^{\mathrm{red}})}$ extends locally uniquely to a locally constant function $c:U\rightarrow\mathbb{C}$ defined near $i(x)$. Writing $f_0=f-c$, we have $f_0|_{i(X^{\mathrm{red}})}=0$ near $i(x)$. The local section s now splits uniquely as $s=s_0+t$ with $\iota_{R,U}(s_0)=f_0+(I'_{R,U})^2$ and $\iota_{R,U}(t)=c+(I'_{R,U})^2$. But s_0 is a local section of $\mathcal{S}_X^0$ as $f_0|_{i(X^{\mathrm{red}})}=0$, and t a local section of $\mathbb{C}_X$ as c is locally constant. Hence $\mathcal{S}_X=\mathcal{S}_X^0\oplus\mathbb{C}_X$, proving (a).

Part (b) involves the *cotangent complex* $\mathbb{L}_X\in D(\mathrm{qcoh}(X))$. For background on cotangent complexes, see Illusie [10], [11, §1]. We need only two facts: that $h^0(\mathbb{L}_X)\cong T^*X$, and if $R\subseteq X$ is open and $i:R\hookrightarrow U$ is a closed embedding of R into a complex manifold U, then the truncation $\tau_{\geqslant -1}(\mathbb{L}_X)$ satisfies

$$\tau_{\geqslant -1}(\mathbb{L}_X)|_R\cong\bigl[I_{R,U}/I_{R,U}^2\xrightarrow{\gamma}i^*(T^*U)\bigr] \tag{3.9}$$

in $D(\mathrm{qcoh}(R))$, where $I_{R,U}/I_{R,U}^2$ is in degree -1 and $i^*(T^*U)$ in degree 0, and the morphism γ maps $\gamma:f+I_{R,U}^2\mapsto i^*(\mathrm{d}f)$.

Consider the diagram of sheaves on R:

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & & \\
0 \arrow[r] & h^{-1}(\mathbb{L}_X)|_R \arrow[r] \arrow[d, dotted, "\alpha_X|_R"] & I_{R,U}/I_{R,U}^2 \arrow[r, "\gamma"] \arrow[d] & i^*(T^*U) \arrow[r] \arrow[d, "\cong"] & \begin{array}{c} h^0(\mathbb{L}_X)|_R \\ \cong T^*X|_R \end{array} \arrow[r] & 0 \\
0 \arrow[r] & \mathcal{S}_X|_R \arrow[r, "\iota_{R,U}"] \arrow[d, dotted, "\beta_X|_R"] & \dfrac{i^{-1}(\mathcal{O}_U)}{I_{R,U}^2} \arrow[r, "\mathrm{d}"] \arrow[d, "\pi"] & \dfrac{i^{-1}(T^*U)}{I_{R,U}\cdot i^{-1}(T^*U)} \arrow[d] & & \\
0 \arrow[r] & \mathcal{O}_X|_R \arrow[r, "\delta", "\cong"'] \arrow[d, "\mathrm{d}"] & \dfrac{i^{-1}(\mathcal{O}_U)}{I_{R,U}} \arrow[r] \arrow[d] & 0 \arrow[d] & & \\
 & T^*X|_R & 0 & 0. & &
\end{tikzcd} \tag{3.10}$$

Here the first row is exact by (3.9), the second row by (2.2), and the third row and second and third columns are obviously exact. Also the middle two squares commute by definition of γ in (3.9). Properties of exact sequences now imply there are unique morphisms $\alpha_X|_R, \beta_X|_R$ as shown making (3.10) commute. Using functoriality of the isomorphism (3.9), one can show that these $\alpha_X|_R, \beta_X|_R$ are independent of R, U, i locally, and so glue on an open cover to give unique global morphisms α_X, β_X in (2.6), noting that the first column of (3.10) is the restriction of (2.6) to R.

Since taking kernels of morphisms of sheaves of $\mathbb{C}$-vector spaces is a left exact functor from the category of morphisms of such sheaves to the category of such sheaves, equation (3.10) also implies that the first column of (3.10) is exact at $h^{-1}(\mathbb{L}_X)|_R$ and $\mathcal{S}_X|_R$. To prove it is exact at $\mathcal{O}_X|_R$, we work at the level of stalks. Let $x \in R \subseteq X$, and write $\mathcal{S}_{X,x}, \mathcal{O}_{X,x}, T^*X_x, \ldots$ for the stalks of $\mathcal{S}_X, \mathcal{O}_X, T^*X, \ldots$ at $x \in X$, so that $\mathcal{S}_{X,x}, \ldots$ are $\mathbb{C}$-vector spaces whose elements are germs at x of sections of $\mathcal{S}_X, \ldots$. A sequence of sheaves of $\mathbb{C}$-vector spaces on R is exact if and only if it is exact on stalks at every $x \in R$.

Let $\eta \in \mathcal{O}_{X,x}$ with $\mathrm{d}\eta = 0 \in T^*X_x$. Then $\delta_x(\eta) \in i^{-1}(\mathcal{O}_U)_x / I_{R,U,x}$, so we may write $\delta_x(\eta) = \zeta + I_{R,U,x}$ for some $\zeta \in i^{-1}(\mathcal{O}_U)_x$. The exact sequences in (3.10) induce an isomorphism

$$T^*X_x \cong i^{-1}(T^*U)_x / \bigl(I_{R,U,x} \cdot i^{-1}(T^*U)_x + \mathrm{d}(I_{R,U,x})\bigr),$$

which identifies $\mathrm{d}\eta \in T^*X_x$ with $\mathrm{d}\zeta + \bigl(I_{R,U,x} \cdot i^{-1}(T^*U)_x + \mathrm{d}(I_{R,U,x})\bigr)$, so $\mathrm{d}\zeta \in I_{R,U,x} \cdot i^{-1}(T^*U)_x + \mathrm{d}(I_{R,U,x}) \subseteq i^{-1}(T^*U)_x$ as $\mathrm{d}\eta = 0$. As ζ is unique up to addition of an element of $I_{R,U,x}$, by changing our choice of ζ we can eliminate the $\mathrm{d}(I_{R,U,x})$ component in $\mathrm{d}\zeta$, so that $\mathrm{d}\zeta \in I_{R,U,x} \cdot i^{-1}(T^*U)_x$. Then $\zeta + I_{R,U,x}^2$ lies in $i^{-1}(\mathcal{O}_U)_x / I_{R,U,x}^2$ with $\mathrm{d}(\zeta + I_{R,U,x}^2) = 0$ in $i^{-1}(T^*U)_x / I_{R,U,x} \cdot i^{-1}(T^*U)_x$. Hence by the exactness of the middle row of (3.10) in stalks at x, there exists a unique $\theta \in \mathcal{S}_{X,x}$ with $\iota_{R,U,x}(\theta) = \zeta + I_{R,U,x}^2$. Therefore

$$\delta_x \circ \beta_{X,x}(\theta) = \pi_x \circ \iota_{R,U,x}(\theta) = \pi_x(\zeta + I_{R,U,x}^2) = \zeta + I_{R,U,x} = \delta_x(\eta),$$

as the bottom left square in (3.10) commutes.

Since δ is an isomorphism, this forces $\beta_{X,x}(\theta) = \eta$. Similarly, if $\theta \in \mathcal{S}_{X,x}$ then $\mathrm{d} \circ \beta_{X,x}(\theta) = 0$. Thus the first column of (3.10) is exact on stalks at $\mathcal{O}_X|_R$, so it is exact. As such open $R \subseteq X$ cover X, we see that (2.6) is exact. Exactness of (2.7) follows from (2.5)–(2.6). This proves Theorem 2.1(b).

For part (c), consider the morphism of sheaves on R

$$\mathrm{d} : i^{-1}(\mathcal{O}_U)/I_{R,U}^2 \longrightarrow i^{-1}(T^*U)/\mathrm{d}(I_{R,U}^2).$$

Composing this with $\iota_{R,U}$ maps $\mathcal{S}_X|_R \to i^{-1}(T^*U)/\mathrm{d}(I_{R,U}^2)$. From (2.2), the image of $\mathrm{d} \circ \iota_{R,U}$ lies in $\bigl(I_{R,U} \cdot i^{-1}(T^*U)/\mathrm{d}(I_{R,U}^2)\bigr) \subseteq i^{-1}(T^*U)/\mathrm{d}(I_{R,U}^2)$. Also, as $\mathrm{d}^2 = 0$, the image lies in $\mathrm{Ker}\,\mathrm{d} + \mathrm{d}(I_{R,U}^2)$. This defines a morphism

$$\mathrm{d} \circ \iota_{R,U} : \mathcal{S}_X|_R \longrightarrow \frac{\mathrm{Ker}\bigl(\mathrm{d} : I_{R,U} \cdot i^{-1}(T^*U) \longrightarrow i^{-1}(\Lambda^2 T^*U)\bigr)}{\mathrm{Im}\bigl(\mathrm{d} : I_{R,U}^2 \longrightarrow I_{R,U} \cdot i^{-1}(T^*U)\bigr)}, \tag{3.11}$$

where the right hand side is the cohomology of (2.8).

In the splitting $\mathcal{S}_X|_R = \mathcal{S}_X^0|_R \oplus \mathbb{C}_X|_R$, clearly $\mathbb{C}_X|_R$ lies in the kernel of $\mathrm{d} \circ \iota_{R,U}$ in (3.11), since d of a locally constant function is zero. We claim that the restriction of (3.11) to $\mathcal{S}_X^0|_R \subset \mathcal{S}_R$ is an isomorphism. As for part (b), it is enough to prove this on the stalks at each $x \in R$. By the Poincaré Lemma on the complex manifold U, written using morphisms $\mathrm{d} : \Lambda^k T^*U_{i(x)} \to \Lambda^{k+1} T^*U_{i(x)}$ of stalks at $i(x)$, and pulled back to X using i, we have an exact sequence

$$\begin{tikzcd} 0 \arrow[r] & \mathbb{C} \arrow[r, "1"] & i^{-1}(\mathcal{O}_X)_x \arrow[r, "\mathrm{d}"] & i^{-1}(T^*U)_x \arrow[r, "\mathrm{d}"] & i^{-1}(\Lambda^2 T^*U)_x. \end{tikzcd} \tag{3.12}$$

Let ϕ lie in the stalk at x of the r.h.s. of (3.11). Then $\phi = \psi + \mathrm{d}(I_{R,U,x}^2)$, where $\psi \in I_{R,U,x} \cdot i^{-1}(T^*U)_x \subseteq i^{-1}(T^*U)_x$ with $\mathrm{d}\psi = 0$ in $i^{-1}(\Lambda^2 T^*U)_x$. By exactness of (3.12), we may write $\psi = \mathrm{d}\zeta$ for $\zeta \in i^{-1}(\mathcal{O}_U)_x$, where ζ is unique up to addition of a constant $1(c)$ for $c \in \mathbb{C}$. We fix ζ uniquely by requiring that $\zeta(i(x)) = 0$. Then $\zeta + I_{R,U,x}^2$ lies in $i^{-1}(\mathcal{O}_U)_x/I_{R,U,x}^2$ with $\mathrm{d}(\zeta + I_{R,U,x}^2) = 0$ in $i^{-1}(T^*U)_x/(I_{R,U,x} \cdot i^{-1}(T^*U)_x)$, since $\mathrm{d}\zeta = \psi \in I_{R,U,x} \cdot i^{-1}(T^*U)_x$. Hence $\zeta + I_{R,U,x}^2 = \iota_{R,U,x}(\theta)$ for unique $\theta \in \mathcal{S}_{X,x}$ as (2.2) is exact, and $\zeta(i(x)) = 0$ implies that $\theta \in \mathcal{S}_{X,x}^0$. Therefore $\mathrm{d} \circ \iota_{R,U}|_{\mathcal{S}_X|_R}$ is an isomorphism on stalks, and so is an isomorphism, proving Theorem 2.1(c).

3.3 Modifications to the proof for the algebraic case

Finally we explain how to modify §3.1–§3.2 to work with $\mathbb{K}$-schemes in algebraic geometry, rather than complex analytic spaces and complex manifolds.

In §3.1 we replace complex analytic spaces X, R by $\mathbb{K}$-schemes, and complex manifolds U by smooth $\mathbb{K}$-schemes, and open subsets $R \subseteq X$, $U' \subseteq U$, etc., are taken to be open in the Zariski topology. Then the proofs work in the algebraic case without modification, with the exception of Lemma 3.2, which is false: one can write down examples of $\mathbb{K}$-schemes X with embeddings $i : X \hookrightarrow U$,

$j : X \hookrightarrow V$ into smooth $\mathbb{K}$-schemes U, V, such that for Zariski or étale open $\emptyset \neq U' \subseteq U$, there exist no morphisms $\Phi : U' \to V$ with $\Phi \circ i' = j|_{U'}$.

Here are two different ways to fix this:

(A) The analogue of Lemma 3.2 is true if we require that the smooth $\mathbb{K}$-scheme V is isomorphic to a Zariski open subset of an affine space $\mathbb{A}^n$, as then we can take the coordinates $(z_1, \ldots, z_n)$ in the proof of Lemma 3.2 to be the embedding $V \hookrightarrow \mathbb{A}^n$.

(B) Suppose $R, S \subseteq X$ are Zariski open and $i : R \hookrightarrow U$, $j : S \hookrightarrow V$ are closed embeddings into smooth $\mathbb{K}$-schemes U, V. Write $\hat{U}, \hat{V}$ for the formal completions of U, V along $i(R), j(S)$, with inclusions $\hat{\imath} : R \hookrightarrow \hat{U}$, $\hat{\jmath} : S \hookrightarrow \hat{V}$. Then one can prove the following formal analogue of Lemma 3.2: for each $x \in R \cap S \subseteq X$, there exists a Zariski open $\hat{U}' \subseteq \hat{U}$ with $x \in R' \subseteq R \cap S$, and a morphism of formal $\mathbb{K}$-schemes $\hat{\Phi} : \hat{U}' \to \hat{V}$ with $\hat{\Phi} \circ \hat{\imath}' = \hat{\jmath}|_{R'}$, where $R' = \hat{\imath}^{-1}(\hat{U}')$ and $\hat{\imath}' = \hat{\imath}|_{R'}$.

Using approach (A), the whole of §3.1 works provided we restrict to triples (R, U, i) with U isomorphic to a Zariski open in $\mathbb{A}^n$. In particular, we can construct the sheaf $\mathcal{S}_X$ using a family $\{(R_a, U_a, i_a) : a \in A\}$ with $U_a \subseteq \mathbb{A}^{n_a}$ Zariski open. Note that immediately after the proof of Lemma 3.3, Zariski locally near each $x \in X$ there exists an embedding $i : X \hookrightarrow U$ with $U \subseteq \mathbb{A}^n$.

The disadvantage of this is that it proves Theorem 2.1(i),(ii) only for U, V Zariski open in affine spaces $\mathbb{A}^n$. To prove them for U, V general smooth $\mathbb{K}$-schemes, we have to do some more work.

To prove the algebraic version of §3.1 using (B), observe that (2.2)–(2.3) defining $\mathcal{S}_X$ depend only on the formal completion $\hat{U}$ of U along $i(R)$, since

$$\frac{i^{-1}(\mathcal{O}_U)}{I_{R,U}^2} \cong \frac{\hat{\imath}^{-1}(\mathcal{O}_{\hat{U}})}{I_{R,\hat{U}}^2} \quad \text{and} \quad \frac{i^{-1}(T^*U)}{I_{R,U} \cdot i^{-1}(T^*U)} \cong \frac{\hat{\imath}^{-1}(T^*\hat{U})}{I_{R,\hat{U}} \cdot \hat{\imath}^{-1}(T^*\hat{U})}.$$

Thus we may replace U, V by $\hat{U}, \hat{V}$ throughout the proof. For triples (R, U, i), (S, V, j) with U, V smooth $\mathbb{K}$-schemes and $R \subseteq S$, a morphism $\Phi : U \to V$ with $\Phi \circ i = j|_R$ induces a morphism $\hat{\Phi} : \hat{U} \to \hat{V}$ with $\hat{\Phi} \circ \hat{\imath} = \hat{\jmath}|_R$. However, the converse is false: there may be $\hat{\Phi} : \hat{U} \to \hat{V}$ with $\hat{\Phi} \circ \hat{\imath} = \hat{\jmath}|_R$ which are not induced by any $\Phi : U \to V$ Zariski or étale locally on U. Because there are more formal morphisms $\hat{\Phi}$ than morphisms Φ, the formal analogue of Lemma 3.2 holds, which makes the proof work.

For the extension of §3.2 to $\mathbb{K}$-schemes, the proofs of Theorem 2.1(a),(b) need no modification. But for (c), equation (3.12) is not exact at $i^{-1}(T^*U)_x$ in the algebraic case: algebraic closed 1-forms need not be locally exact, in either the Zariski or the étale topology. For example, the closed 1-form $z^{-1}\mathrm{d}z$ on $\mathbb{A}^1 \setminus \{0\}$ is not algebraically locally exact, since $\log z$ is not an algebraic function.

Because of this, in the algebraic case the morphism from $\mathcal{S}_X^0$ to the cohomology of (2.8) constructed in §3.2 is injective, but generally not surjective. The solution is to modify (2.8), replacing U by the formal completion $\hat{U}$ of U along $i(R)$, as in (2.9). In fact the Poincaré Lemma may not hold on $\hat{U}$ either (consider the case $X = U = \hat{U}$ and $i = \mathrm{id}_X$), but what matters is that one can show that if $\hat{\alpha} \in H^0(I_{R,\hat{U}} \cdot T^*\hat{U}) \subset H^0(T^*\hat{U})$ with $\mathrm{d}\hat{\alpha} = 0$ then $\hat{\alpha} = \mathrm{d}f$ for a unique $f \in H^0(\mathcal{O}_{\hat{U}})$ with $f \circ i|_{X^{\mathrm{red}}} = 0$, that is, the particular class of formal closed 1-forms we are interested in can be integrated to formal functions.

4 Comparing critical charts (R, U, f, i)

Next we prove Theorem 2.13 and Propositions 2.12 and 2.15–2.17 from §2.3.

4.1 Proof of Proposition 2.12

Let (R, U, f, i) be a critical chart on an algebraic d-critical locus (X, s) over $\mathbb{K}$, and $x \in R$. Then U is a smooth $\mathbb{K}$-scheme, so there exist an affine open neighbourhood $\tilde{U}$ of $i(x)$ in U and a closed embedding of $\mathbb{K}$-schemes $\tilde{\Phi} : \tilde{U} \hookrightarrow \mathbb{A}^n$ for some $n \geqslant 0$. Choose a Zariski open neighbourhood V of $\tilde{\Phi}(i(x))$ in $\mathbb{A}^n$ and étale coordinates $(z_1, \ldots, z_n)$ on V such that $\tilde{\Phi}(\tilde{U}) \cap V$ is the $\mathbb{K}$-subscheme defined by $z_{m+1} = \cdots = z_n = 0$ in V, where $m = \dim U$. Set $U' = \tilde{\Phi}^{-1}(V)$, $R' = S = i^{-1}(U')$, $i' = i|_{R'} : R' \hookrightarrow U'$, $f' = f|_{U'} : U' \to \mathbb{A}^1$, $\Phi = \tilde{\Phi}|_{U'} : U' \hookrightarrow V$, and $j = \Phi \circ i' : S = R' \hookrightarrow V$.

Now $\Phi(U')$ is a closed $\mathbb{K}$-subscheme of V, and $f' \circ \Phi^{-1} : \Phi(U') \to \mathbb{A}^1$ a regular function. Zariski locally on V, we may extend $f' \circ \Phi^{-1}$ from $\Phi(U')$ to V. Thus, making V, U', R', S smaller, there exists a regular $h : V \to \mathbb{A}^1$ with $h \circ \Phi = f' : U' \to \mathbb{A}^1$. Then we may form the partial derivatives $\frac{\partial h}{\partial z_a}, \frac{\partial^2 h}{\partial z_a \partial z_b}$ for $a, b = 1, \ldots, n$. Define $g : V \to \mathbb{A}^1$ by

$$g = h - \textstyle\sum_{a=m+1}^{n} z_a \cdot \frac{\partial h}{\partial z_a} + \frac{1}{2} \sum_{a,b=m+1}^{n} z_a z_b \cdot \frac{\partial^2 h}{\partial z_a \partial z_b} + \sum_{a=m+1}^{n} z_a^2. \tag{4.1}$$

Here we use $\mathrm{char}\,\mathbb{K} \neq 2$ from §1. Since $z_a|_{\Phi(U')} = 0$ for $a = m+1, \ldots, n$, we have $g \circ \Phi = h \circ \Phi = f' : U' \to \mathbb{A}^1$. Also equation (4.1) implies that

$$\tfrac{\partial g}{\partial z_a}|_{\Phi(U')} = 0, \;\; a = m+1, \ldots, n, \;\; \tfrac{\partial^2 g}{\partial z_a \partial z_b}|_{\Phi(U')} = 2\delta_{ab}, \;\; a, b = m+1, \ldots, n. \tag{4.2}$$

Consider the $\mathbb{K}$-subscheme of V defined by the equations $\frac{\partial g}{\partial z_a} = 0$ for $a = m+1, \ldots, n$. The first equation of (4.2) shows this $\mathbb{K}$-subscheme contains $\Phi(U')$, and the second that this $\mathbb{K}$-subscheme coincides with $\Phi(U')$ near $\Phi(U')$ in V. Hence, by making V smaller while keeping U', R', S fixed, we can take this $\mathbb{K}$-subscheme to be $\Phi(U')$. We now claim that

$$\mathrm{Crit}(g) = \Phi(\mathrm{Crit}(f')) = \Phi(i(R')) = j(S) \tag{4.3}$$

as $\mathbb{K}$-subschemes of V. To see this, note that the equations $\frac{\partial g}{\partial z_a} = 0$ for $a = 1, \ldots, m$ defining $\mathrm{Crit}(g)$ divide into $\frac{\partial g}{\partial z_a} = 0$ for $a = m+1, \ldots, n$, which define the smooth $\mathbb{K}$-subscheme $\Phi(U')$ in V, and $\frac{\partial g}{\partial z_a} = 0$ for $a = 1, \ldots, m$, which under $\Phi(U') \cong U'$ correspond to $\mathrm{d}f' = 0$ as $g \circ \Phi = f'$.

As (R', U', f', i') is a critical chart on (X, s) we have $\iota_{R',U'}(s|_{R'}) = i'^{-1}(f') + I_{R',U'}^2$. Applying Theorem 2.1(ii) with (R', U', i) in place of (R, U, i) and using

$g \circ \Phi = f'$, $j = \Phi \circ i'$ and $R' = S$ shows that $\iota_{S,V}(s|_S) = j^{-1}(g) + I_{S,V}^2$. Together with (4.3), this implies that (S, V, g, j) is a critical chart, and $\Phi : U' \hookrightarrow V$ an embedding with $j = \Phi \circ i'$, $g \circ \Phi = f'$ show $\Phi : (R, U, f, i) \hookrightarrow (S, V, g, j)$ is an embedding of critical charts. This proves Proposition 2.12.

4.2 Proof of Theorem 2.13

We begin with the complex analytic case. Let (X, s) be a complex analytic d-critical locus, $(R, U, f, i), (S, V, g, j)$ be critical charts on (X, s), and $x \in R \cap S$. Then $i(R \cap S)$ is a locally closed complex analytic subspace of U, and $j \circ i^{-1}|_{i(R\cap S)} : i(R \cap S) \to V$ a morphism to a complex manifold V. So we can extend $j \circ i^{-1}$ locally to a holomorphic map $U \to V$. That is, we can choose an open neighbourhood U' of $i(x)$ in U with $R' := i^{-1}(U') \subseteq R \cap S$, and a holomorphic map $\Theta : U' \to V$ such that $\Theta \circ i' = j|_{R'} : R' \to V$, for $i' := i|_{R'}$.

Theorem 2.1(ii) with $(R', U', i'), \Theta$ in place of $(R, U, i), \Phi$ gives a commutative diagram (2.3). Applying this to $s|_{R'}$ shows that

$$\iota_{R',U'}(s|_{R'}) = i'^{-1}(\Theta^\sharp)\bigl[\iota_{S,V}(s|_S)\bigr]. \tag{4.4}$$

Write $I'_{R',U'} \subseteq \mathcal{O}_{U'}$ for the ideal vanishing on $i(R') \subseteq U'$, and $f' = f|_{U'}$. Then

$$\begin{aligned}
i'^{-1}\bigl[f' + (I'_{R',U'})^2\bigr] &= i'^{-1}(f') + I_{R',U'}^2 = \iota_{R',U'}(s|_{R'}) = i'^{-1}(\Theta^\sharp)\bigl[\iota_{S,V}(s|_S)\bigr] \\
&= i'^{-1}(\Theta^\sharp)\bigl[j^{-1}(g) + I_{S,V}^2\bigr] = i'^{-1}(\Theta^\sharp)\bigl[i'^{-1} \circ \Theta^{-1}(g) + I_{S,V}^2\bigr] \\
&= i'^{-1}\bigl[\Theta^\sharp\bigl(\Theta^{-1}(g)\bigr)\bigr] + I_{R',U'}^2 = i'^{-1}\bigl[g \circ \Theta + (I'_{R',U'})^2\bigr],
\end{aligned} \tag{4.5}$$

using $i'^{-1}(I'_{R',U'}) = I_{R',U'}$ in the first and seventh steps, $\iota_{R',U'}(s|_{R'}) = i'^{-1}(f') + I_{R',U'}^2$ in the second, (4.4) in the third, $\iota_{S,V}(s|_S) = j^{-1}(g) + I_{S,V}^2$ in the fourth, $\Theta \circ i' = j|_{R'}$ in the fifth, and $i'^{-1}(\Theta^\sharp)(I_{S,V}) = I_{R',U'}$ in the sixth.

Equation (4.5) implies that $f' - g \circ \Theta \in (I'_{R',U'})^2$. Therefore, making U', R' smaller if necessary, we can choose holomorphic functions $r_a, s_a : U' \to \mathbb{C}$ for $a = 1, \ldots, n$, some $n \geqslant 0$, such that $r_a, s_a \in H^0(I'_{R',U'})$ for $a = 1, \ldots, n$ and

$$f' = g \circ \Theta + r_1 s_1 + \cdots + r_n s_n : U' \longrightarrow \mathbb{C}. \tag{4.6}$$

Define $W = V \times \mathbb{C}^{2n}$, $T = S$, $k = j \times (0, \ldots, 0) : T = S \hookrightarrow V \times \mathbb{C}^{2n} = W$, $(S', V', g', j') = (S, V, g, j)$, and $\Phi : U' \to W$, $\Psi : V' \to W$, $h : W \to \mathbb{C}$ by

$$\begin{gathered}
\Phi(u) = \bigl(\Theta(u), (r_1(u), \ldots, r_n(u), s_1(u), \ldots, s_n(u))\bigr), \quad \Psi(v) = \bigl(v, (0, \ldots, 0)\bigr), \\
\text{and} \quad h\bigl(v, (y_1, \ldots, y_n, z_1, \ldots, z_n)\bigr) = g(v) + y_1 z_1 + \cdots + y_n z_n.
\end{gathered} \tag{4.7}$$

By increasing n and adding extra functions r_a, s_a with $s_a = 0$, we can suppose Φ is an embedding near $i(x) \in U'$. So making U', R' smaller we can take $\Phi : U' \hookrightarrow W$ to be an embedding.

Since (S, V, g, j) is a critical chart, (T, W, h, k) is one too. Also $\Phi : U' \hookrightarrow W$, $\Psi : V' \hookrightarrow W$ are embeddings, and $f' = h \circ \Phi$, $g' = h \circ \Psi$, $\Phi \circ i' = k|_{R'}$, $\Psi \circ j' = k|_{S'}$ follow from (4.6)–(4.7) and $\Theta \circ i' = j|_{R'}$. Therefore $\Phi : (R', U', f', i') \hookrightarrow$

(T,W,h,k), $\Psi:(S',V',g',j')\hookrightarrow(T,W,h,k)$ are embeddings of critical charts with $x\in R'\cap S'$, proving Theorem 2.13 in the complex analytic case.

For the algebraic case, with (X,s) an algebraic d-critical locus over $\mathbb{K}$, and $(R,U,f,i),(S,V,g,j)$ critical charts, we would like to follow the above method, but there is a problem with the first step: if V is a general smooth $\mathbb{K}$-scheme, we may not be able to choose a Zariski open $i(x)\in U'\subseteq U$ and a morphism $\Theta:U'\rightarrow V$ such that $\Theta\circ i'=j|_{R'}:R'\rightarrow V$. However, this is valid if V is Zariski open in an affine space $\mathbb{A}^m$.

So we modify the method above as follows: first we apply Proposition 2.12, proved in §4.1, to get a subchart $(S',V',g',j')\subseteq(S,V,g,j)$ with $x\in S'$ and an embedding $\Xi:(S',V',g',j')\hookrightarrow(\tilde S,\tilde V,\tilde g,\tilde\jmath)$ for $(\tilde S,\tilde V,\tilde g,\tilde\jmath)$ a critical chart on (X,s) with $\tilde V\subseteq\mathbb{A}^m$ Zariski open. Thus we may choose Zariski open $i(x)\in U'\subseteq U$ and a morphism $\Theta:U'\rightarrow\tilde V$ with $\Theta\circ i'=\tilde\jmath|_{R'}:R'\rightarrow W$. Then we follow the rest of the proof above with $(\tilde S,\tilde V,\tilde g,\tilde\jmath),\mathbb{A}^n$ in place of $(S,V,g,j),\mathbb{C}^n$, so that $W=\tilde V\times\mathbb{A}^{2n}$, except that we define $\Psi:V'\hookrightarrow W$ by $\Psi(v)=\bigl(\Xi(v),(0,\ldots,0)\bigr)$. We leave the details to the reader.

4.3 Proof of Proposition 2.15

Choose holomorphic coordinates $(\dot y_1,\ldots,\dot y_m,\dot z_1,\ldots,\dot z_n)$ on an open neighbourhood $\dot V$ of $j(x)$ in V, where $\dim U=m$ and $\dim V=m+n$, such that $j(x)=(0,\ldots,0)$ and $\Phi(U)\cap\dot V$ is the submanifold $\dot z_1=\cdots=\dot z_n=0$ in $\dot V$. Set $\dot U=\Phi^{-1}(\dot V)$ and $\dot x_a=\dot y_a\circ\Phi|_{\dot U}$ for $a=1,\ldots,m$. Then $\dot U$ is an open neighbourhood of $i(x)$ in U, and $(\dot x_1,\ldots,\dot x_m)$ are holomorphic coordinates on $\dot U$ with $i(x)=(0,\ldots,0)$. Write $f|_{\dot U}=\dot f(\dot x_1,\ldots,\dot x_m)$ and $g|_{\dot V}=\dot g(\dot y_1,\ldots,\dot y_m,\dot z_1,\ldots,\dot z_n)$ as functions of these coordinates, so that $f=g\circ\Phi$ implies that $\dot f(\dot y_1,\ldots,\dot y_m)=\dot g(\dot y_1,\ldots,\dot y_m,0,\ldots,0)$ for $(\dot y_1,\ldots,\dot y_m)\in\dot U$.

Then the ideal $I'_{R,U}=I_{(\mathrm{d}f)}$ is on $\dot U$ the ideal of holomorphic functions in $(\dot x_1,\ldots,\dot x_m)$ generated by $\frac{\partial\dot f}{\partial\dot x_a}$ for $a=1,\ldots,m$, and the ideal $I'_{S,V}=I_{(\mathrm{d}g)}$ is on $\dot V$ the ideal of holomorphic functions in $(\dot y_1,\ldots,\dot y_m,\dot z_1,\ldots,\dot z_n)$ generated by $\frac{\partial\dot g}{\partial\dot y_a}$ for $a=1,\ldots,m$ and $\frac{\partial\dot g}{\partial\dot z_b}$ for $b=1,\ldots,n$. Since Φ maps U to $\dot z_1=\cdots=\dot z_n=0$ and $i(R)$ to $j(R)\subseteq j(S)$, we have $I_{(\mathrm{d}f)}\cong I_{(\mathrm{d}g)}|_{\dot z_1=\cdots=\dot z_n=0}$, that is,

$$\bigl(\tfrac{\partial\dot f}{\partial\dot x_a}(\dot y_1,\ldots,\dot y_m):a=1,\ldots,m\bigr)=$$
$$\bigl(\tfrac{\partial\dot g}{\partial\dot y_a}(\dot y_1,\ldots,\dot y_m,0,\ldots,0):a{=}1,\ldots,m,\;\;\tfrac{\partial\dot g}{\partial\dot z_b}(\dot y_1,\ldots,\dot y_m,0,\ldots,0):b{=}1,\ldots,n\bigr).$$

As $\frac{\partial\dot g}{\partial\dot y_a}(\dot y_1,\ldots,\dot y_m,0,\ldots,0)=\frac{\partial\dot f}{\partial\dot x_a}(\dot y_1,\ldots,\dot y_m)$, this holds provided each $\frac{\partial\dot g}{\partial\dot z_b}(\dot y_1,\ldots,\dot y_m,0,\ldots,0)$ lies in $\bigl(\frac{\partial\dot g}{\partial\dot y_a}(\dot y_1,\ldots,\dot y_m,0,\ldots,0):a=1,\ldots,m\bigr)$. Thus, making $\dot U,\dot V$ smaller if necessary, we can suppose there exist holomorphic functions $A_{ab}(\dot y_1,\ldots,\dot y_m)$ on $\dot U$ for $a=1,\ldots,m$, $b=1,\ldots,n$ such that for each b

$$\tfrac{\partial\dot g}{\partial\dot z_b}(\dot y_1,\ldots,\dot y_m,0,\ldots,0)=\textstyle\sum_{a=1}^m A_{ab}(\dot y_1,\ldots,\dot y_m)\cdot\tfrac{\partial\dot g}{\partial\dot y_a}(\dot y_1,\ldots,\dot y_m,0,\ldots,0).$$

Define holomorphic coordinates $(\tilde y_1,\ldots,\tilde y_m,\tilde z_1,\ldots,\tilde z_n)$ on an open neighbourhood $\tilde V$ of $j(x)$ in $\dot V$ by $\tilde y_a=\dot y_a+\sum_{b=1}^n A_{ab}(\dot y_1,\ldots,\dot y_m)\,\dot z_b$ and $\tilde z_b=\dot z_b$.

Here $\tilde{y}_a, \tilde{z}_b$ are defined on all of $\dot{V}$, but they need only be a coordinate system near $j(x)$ in $\dot{V}$ where $\dot{z}_1, \ldots, \dot{z}_n$ are small, so we shrink $\dot{V}$ to $\tilde{V} \subseteq \dot{V}$. We also write $\tilde{U} = \Phi^{-1}(\tilde{V}) \subseteq \dot{U}$ and $\tilde{x}_a = \dot{x}_a|_{\tilde{U}}$. Then $\tilde{y}_a \circ \Phi|_{\tilde{U}} = \tilde{x}_a$ as $\dot{z}_b \circ \Phi|_{\tilde{U}} = 0$. Making $\tilde{V}$ smaller if necessary we can suppose that if $(\tilde{y}_1, \ldots, \tilde{y}_m, \tilde{z}_1, \ldots, \tilde{z}_n) \in \tilde{V}$ then $(\tilde{y}_1, \ldots, \tilde{y}_m) \in \dot{U}$, using the coordinates $(\dot{x}_1, \ldots, \dot{x}_m)$ on $\dot{U}$.

Write $g|_{\tilde{V}} = \tilde{g}(\tilde{y}_1, \ldots, \tilde{y}_m, \tilde{z}_1, \ldots, \tilde{z}_n)$ using these coordinates. Then

$$\begin{aligned}
&\tfrac{\partial \tilde{g}}{\partial \tilde{z}_b}(\tilde{y}_1, \ldots, \tilde{y}_m, 0, \ldots, 0) \\
&= \textstyle\sum_{a=1}^m \tfrac{\partial \dot{g}}{\partial \dot{y}_a}(\dot{y}_1, \ldots, \dot{y}_m, 0, \ldots, 0) \cdot \tfrac{\partial \dot{y}_a}{\partial \tilde{z}_b} + \textstyle\sum_{c=1}^n \tfrac{\partial \dot{g}}{\partial \dot{z}_c}(\dot{y}_1, \ldots, \dot{y}_m, 0, \ldots, 0) \cdot \tfrac{\partial \dot{z}_c}{\partial \tilde{z}_b} \\
&= \textstyle\sum_{a=1}^m \tfrac{\partial \dot{g}}{\partial \dot{y}_a}(\dot{y}_1, \ldots, \dot{y}_m, 0) \cdot (-A_{ab}(\dot{y}_1, \ldots, \dot{y}_m)) \\
&\quad + \textstyle\sum_{c=1}^n \bigl(\textstyle\sum_{a=1}^m A_{ac}(\dot{y}_1, \ldots, \dot{y}_m) \cdot \tfrac{\partial \dot{g}}{\partial \dot{y}_a}(\dot{y}_1, \ldots, \dot{y}_m, 0, \ldots, 0)\bigr) \cdot \delta_{bc} = 0.
\end{aligned}$$

So $\tilde{g}(\tilde{y}_1, \ldots, \tilde{y}_m, 0, \ldots, 0) = \dot{f}(\tilde{y}_1, \ldots, \tilde{y}_m)$ and $\frac{\partial \tilde{g}}{\partial \tilde{z}_b}(\tilde{y}_1, \ldots, \tilde{y}_m, 0, \ldots, 0) = 0$ for $b = 1, \ldots, n$ in the new coordinates $(\tilde{y}_1, \ldots, \tilde{y}_m, \tilde{z}_1, \ldots, \tilde{z}_n)$.

Consider the holomorphic function $h : \tilde{V} \to \mathbb{C}$ given by

$$h(\tilde{y}_1, \ldots, \tilde{y}_m, \tilde{z}_1, \ldots, \tilde{z}_n) = \tilde{g}(\tilde{y}_1, \ldots, \tilde{y}_m, \tilde{z}_1, \ldots, \tilde{z}_n) - \dot{f}(\tilde{y}_1, \ldots, \tilde{y}_m). \tag{4.8}$$

It satisfies $h(\tilde{y}_1, \ldots, \tilde{y}_m, 0, \ldots, 0) = 0$ and $\frac{\partial h}{\partial \tilde{z}_b}(\tilde{y}_1, \ldots, \tilde{y}_m, 0, \ldots, 0) = 0$ for all $b = 1, \ldots, n$ and $(\tilde{y}_1, \ldots, \tilde{y}_m) \in \tilde{U}$, so h lies in the ideal $(\tilde{z}_1, \ldots, \tilde{z}_n)^2$. Thus, making $\tilde{V}, \tilde{U}$ smaller if necessary, we may write

$$h(\tilde{y}_1, \ldots, \tilde{y}_m, \tilde{z}_1, \ldots, \tilde{z}_n) = \textstyle\sum_{b,c=1}^n \tilde{z}_b \tilde{z}_c \, Q_{bc}(\tilde{y}_1, \ldots, \tilde{y}_m, \tilde{z}_1, \ldots, \tilde{z}_n), \tag{4.9}$$

for some (nonunique) holomorphic functions $Q_{bc} : \tilde{V} \to \mathbb{C}$ with $Q_{bc} = Q_{cb}$.

Now $\operatorname{Crit}(\tilde{g}) = \operatorname{Crit}(g) \cap \tilde{V} = j(S) \cap \tilde{V}$ lies in $\Phi(\tilde{U}) = \{\tilde{z}_1 = \cdots = \tilde{z}_n = 0\}$ as complex analytic subspaces of $\tilde{V}$. Therefore $\tilde{z}_1, \ldots, \tilde{z}_n$ lie in the ideal $I_{(\mathrm{d}\tilde{g})}$ generated by $\frac{\partial \tilde{g}}{\partial \tilde{y}_a}, \frac{\partial \tilde{g}}{\partial \tilde{z}_b}$, so making $\tilde{U}, \tilde{V}$ smaller, for each $d = 1, \ldots, n$ there exist holomorphic functions $B_{ad}(\tilde{y}_1, \ldots, \tilde{y}_m, \tilde{z}_1, \ldots, \tilde{z}_n), C_{bd}(\tilde{y}_1, \ldots, \tilde{y}_m, \tilde{z}_1, \ldots, \tilde{z}_n)$ on $\tilde{V}$ such that using (4.8)–(4.9) we have

$$\begin{aligned}
\tilde{z}_d &= \sum_{a=1}^m B_{ad} \cdot \tfrac{\partial \tilde{g}}{\partial \tilde{y}_a} + \sum_{b=1}^n C_{bd} \cdot \tfrac{\partial \tilde{g}}{\partial \tilde{z}_b} \\
&= \sum_{a=1}^m B_{ad} \cdot \Bigl[\tfrac{\partial \dot{f}}{\partial \dot{x}_a}(\tilde{y}_1, \ldots, \tilde{y}_m) + \sum_{a=1}^m \sum_{b,c=1}^n \tilde{z}_b \tilde{z}_c \tfrac{\partial Q_{bc}}{\partial \tilde{y}_a}(\tilde{y}_1, \ldots, \tilde{y}_m, \tilde{z}_1, \ldots, \tilde{z}_n)\Bigr] \\
&\quad + 2 \sum_{b,c=1}^n C_{bd} \cdot \tilde{z}_c Q_{bc}(\tilde{y}_1, \ldots, \tilde{y}_m, \tilde{z}_1, \ldots, \tilde{z}_n) \\
&\quad + \sum_{b,c,e=1}^n C_{ed} \cdot \tilde{z}_b \tilde{z}_c \tfrac{\partial Q_{bc}}{\partial \tilde{z}_e}(\tilde{y}_1, \ldots, \tilde{y}_m, \tilde{z}_1, \ldots, \tilde{z}_n).
\end{aligned} \tag{4.10}$$

Apply $\frac{\partial}{\partial \tilde{z}_c}$ to (4.10) and restrict it to the point $(0, \ldots, 0) = j(x)$, noting that $\frac{\partial \dot{f}}{\partial \dot{x}_a}(0, \ldots, 0) = 0$. This yields

$$\delta_{cd} = 2 \textstyle\sum_{b=1}^n C_{bd}(0, \ldots, 0) \cdot Q_{bc}(0, \ldots, 0), \quad \text{for all } c, d = 1, \ldots, n.$$

Hence the symmetric matrix $\bigl(Q_{bc}(0,\ldots,0)\bigr)_{b,c=1}^n$ is invertible. Thus, by applying an element of $\mathrm{GL}(n,\mathbb{C})$ to the coordinates $(\tilde z_1,\ldots,\tilde z_n)$ we can suppose that $Q_{bc}(0,\ldots,0)=\delta_{bc}$ for $b,c=1,\ldots,n$.

We now define new holomorphic coordinates $(y_1,\ldots,y_m,z_1,\ldots,z_n)$ on an open neighbourhood V' of $j(x)$ in $\tilde V$, and write $g'(y_1,\ldots,y_m,z_1,\ldots,z_n)=g|_{V'}$ for g as a function of these new coordinates and $U'=\Phi^{-1}(V')$, such that:

(a) $y_a=\tilde y_a$ for $a=1,\ldots,m$.

(b) $\frac{\partial z_b}{\partial \tilde z_c}=\delta_{bc}$ at $j(x)$ for $b,c=1,\ldots,n$.

(c) $\Phi(U')$ is the submanifold $z_1=\cdots=z_n=0$ in V'.

(d) $g'(y_1,\ldots,y_m,z_1,\ldots,z_n)=\dot f(y_1,\ldots,y_m)+z_1^2+\cdots+z_n^2$.

We define z_b by reverse induction on $b=n,n-1,\ldots,1$.

For the first step, as $Q_{nn}=1$ at $j(x)=(0,\ldots,0)$, we may restrict to a small open neighbourhood V' of $j(x)$ in $\tilde V$ on which $Q_{nn}(\tilde y_1,\ldots,\tilde y_m,\tilde z_1,\ldots,\tilde z_n)$ is invertible and has a square root $Q_{nn}^{1/2}$. Rewrite (4.9) on V' as

$$\begin{aligned}
&\tilde g(\tilde y_1,\ldots,\tilde y_m,\tilde z_1,\ldots,\tilde z_n)=\dot f(\tilde y_1,\ldots,\tilde y_m)+\textstyle\sum_{b,c=1}^{n-1}\tilde z_b\tilde z_c\Bigl[Q_{bc}-Q_{nn}^{-1}Q_{bn}Q_{cn}\Bigr]\\
&+\Bigl[Q_{nn}^{1/2}\tilde z_n+\textstyle\sum_{b=1}^{n-1}Q_{nn}^{-1/2}Q_{bn}\tilde z_b\Bigr]^2=\dot f(\tilde y_1,\ldots,\tilde y_m)+\textstyle\sum_{b,c=1}^{n-1}\tilde z_b\tilde z_c\hat Q_{bc}+z_n^2,\\
&\text{where }\hat Q_{bc}=Q_{bc}-Q_{nn}^{-1}Q_{bn}Q_{cn},\;\; z_n=Q_{nn}^{1/2}\tilde z_n+\textstyle\sum_{b=1}^{n-1}Q_{nn}^{-1/2}Q_{bn}\tilde z_b.
\end{aligned}\tag{4.11}$$

Note that $\hat Q_{bc}(0,\ldots,0)=\delta_{bc}$ for $b,c=1,\ldots,n-1$, as $Q_{bc}(0,\ldots,0)=\delta_{bc}$. For the second step, making V' smaller so that $\hat Q_{n-1\,n-1}$ is invertible and has a square root $\hat Q_{n-1\,n-1}^{1/2}$ on V', we have

$$\begin{aligned}
\tilde g(\tilde y_1,\ldots,\tilde y_m,\tilde z_1,\ldots,\tilde z_n)&=f(\tilde y_1,\ldots,\tilde y_m)+\textstyle\sum_{b,c=1}^{n-2}\tilde z_b\tilde z_c\check Q_{bc}+z_{n-1}^2+z_n^2,\\
\text{where}\qquad \check Q_{bc}&=\hat Q_{bc}-\hat Q_{n-1\,n-1}^{-1}\hat Q_{b\,n-1}\hat Q_{c\,n-1}\\
\text{and}\qquad z_{n-1}&=\hat Q_{n-1\,n-1}^{1/2}\tilde z_{n-1}+\textstyle\sum_{b=1}^{n-2}\hat Q_{n-1\,n-1}^{-1/2}\hat Q_{b\,n-1}\tilde z_b.
\end{aligned}$$

Continuing in this way, we define $j(x)\in V'\subseteq\tilde V$ and holomorphic functions $y_1,\ldots,y_m,z_1,\ldots,z_n:V'\to\mathbb{C}$ satisfying (a)–(d) above.

Parts (a)–(c) imply that $(y_1,\ldots,y_m,z_1,\ldots,z_n)$ are a coordinate system near $j(x)$ in V', so making V' smaller, we can suppose they are coordinates on V'. Define $\alpha:V'\to U$ and $\beta:V'\to\mathbb{C}^n$ by $\alpha:(y_1,\ldots,y_m,z_1,\ldots,z_n)\mapsto(y_1,\ldots,y_m)$ and $\beta:(y_1,\ldots,y_m,z_1,\ldots,z_n)\mapsto(z_1,\ldots,z_n)$, using coordinates $(y_1,\ldots,y_m,z_1,\ldots,z_n)$ on V' and $(\dot x_1,\ldots,\dot x_m)$ on $\dot U$. Proposition 2.15 then follows from (a),(c),(d) above.

4.4 Proof of Proposition 2.16

We will adapt §4.3 to the algebraic context. The first part, until just before (4.8), works with $i(x)\in\tilde U\subseteq\dot U\subseteq U$ and $j(x)\in\tilde V\subseteq\dot V\subseteq V$ Zariski open and $(\dot x_1,\ldots,\dot x_m):\dot U\to\mathbb{A}^n$, $(\dot y_1,\ldots,\dot y_m,\dot z_1,\ldots,\dot z_n):\dot V\to\mathbb{A}^{m+n}$, $(\tilde y_1,\ldots,\tilde y_m,$

$\tilde{z}_1,\ldots,\tilde{z}_n) : \tilde{V} \to \mathbb{A}^{m+n}$ étale coordinates on $\dot{U}, \dot{V}, \tilde{V}$. Note that f, g are not functions of $\dot{x}_a$ or $\dot{y}_a, \dot{z}_b$ or $\tilde{y}_a, \tilde{z}_b$, except in an étale sense, so we cannot rigorously write $f|_{\dot{U}} = \dot{f}(\dot{x}_1,\ldots,\dot{x}_m)$, and so on. Nonetheless, the partial derivatives $\frac{\partial f}{\partial \dot{x}_a}$, $\frac{\partial g}{\partial \dot{y}_a}, \frac{\partial g}{\partial \dot{z}_b}, \frac{\partial g}{\partial \tilde{y}_a}, \frac{\partial g}{\partial \tilde{z}_b}$ are all well defined on $\dot{U}, \dot{V}, \tilde{V}$.

In this way, we construct étale coordinates $(\tilde{y}_1,\ldots,\tilde{y}_m,\tilde{z}_1,\ldots,\tilde{z}_n)$ on $\tilde{V} \subseteq V$ such that $\Phi(\tilde{U})$ is the smooth $\mathbb{K}$-subscheme $\tilde{z}_1 = \cdots = \tilde{z}_n = 0$ in $\tilde{V}$, and $\frac{\partial g}{\partial \tilde{z}_b}|_{\Phi(\tilde{U})} = 0$ for $b = 1,\ldots,n$. Now (4.8) does not make sense on $\tilde{V}$, since we cannot extend f from $\tilde{U}$ to $\tilde{V}$ as f is not a function of $(\tilde{y}_1,\ldots,\tilde{y}_m)$. Instead, we form the Cartesian square of smooth $\mathbb{K}$-schemes and étale morphisms

$$\begin{tikzcd}
\check{V} = (\tilde{U} \times \mathbb{A}^n) \times_{\mathbb{A}^{m+n}} \tilde{V} \arrow[r, "\check{\jmath}"'] \arrow[d, "\check{\alpha}\times\check{\beta}"] & \tilde{V} \arrow[d, "(\tilde{y}_1,\ldots,\tilde{y}_m,\tilde{z}_1,\ldots,\tilde{z}_n)"'] \\
\tilde{U} \times \mathbb{A}^n \arrow[r, "(\tilde{x}_1,\ldots,\tilde{x}_m)\times\mathrm{id}_{\mathbb{A}^n}"] & \mathbb{A}^{m+n} = \mathbb{A}^m \times \mathbb{A}^n.
\end{tikzcd} \tag{4.12}$$

There is a unique $\check{v} \in \check{V}$ with $\check{\alpha}(\check{v}) = i(x)$, $\check{\beta}(\check{v}) = (0,\ldots,0)$ and $\check{\jmath}(\check{v}) = j(x)$. Then $\check{\alpha} \times \check{\beta}$ and $\check{\jmath}$ are étale, so we can regard $\check{\jmath} : \check{V} \to \tilde{V} \subseteq V$ as an étale open set in V. Define étale coordinates $(\check{y}_1,\ldots,\check{y}_m,\check{z}_1,\ldots,\check{z}_n) : \check{V} \to \mathbb{A}^{m+n}$ by $\check{y}_a = \tilde{y}_a \circ \check{\jmath}$, $\check{z}_b = \tilde{z}_b \circ \check{\jmath}$, and define $\check{f}, \check{g} : \check{V} \to \mathbb{A}^1$ by $\check{f} = f \circ \check{\alpha}$ and $\check{g} = g \circ \check{\jmath}$. The analogue of (4.8) is now $\check{h} = \check{g} - \check{f} : \check{V} \to \mathbb{A}^1$. The previous argument now shows that on the smooth $\mathbb{K}$-subscheme $\check{U} \in \check{V}$ defined by $\check{z}_1 = \cdots = \check{z}_n = 0$ we have $\check{h}|_{\check{U}} = 0$ and $\frac{\partial \check{h}}{\partial \check{z}_b}|_{\check{U}} = 0$ for $b = 1,\ldots,n$. Therefore $\check{h}$ lies in the ideal $(\check{z}_1,\ldots,\check{z}_n)^2$ on $\check{V}$. So making $\tilde{U}, \check{U}, \tilde{V}, \check{V}$ smaller, we may write $\check{h} = \sum_{b,c=1}^n \check{z}_b \check{z}_c Q_{bc}$, the analogue of (4.9), for some $Q_{bc} : \check{V} \to \mathbb{A}^1$ with $Q_{bc} = Q_{cb}$, where the last part requires $\operatorname{char} \mathbb{K} \neq 2$, and $Q_{bc}(0,\ldots,0) = \delta_{bc}$ becomes $Q_{bc}(\check{v}) = \delta_{bc}$.

The last part of §4.3, defining the coordinates $(y_1,\ldots,y_m,z_1,\ldots,z_n)$, involves taking square roots $Q_{nn}^{1/2}, \hat{Q}_{n-1\,n-1}^{-1/2}, \ldots$. These generally will not exist on $\check{V}$ or Zariski open subsets of $\check{V}$, but they will exist on étale open subsets of $\check{V}$, noting that $\mathbb{K}$ is algebraically closed, so that square roots exist in $\mathbb{K}$. So we can construct an étale open neighbourhood $\jmath' : V' \to \check{V}$ of $\check{v}$ in $\check{V}$ on which we define étale coordinates $(y_1,\ldots,y_m,z_1,\ldots,z_n) : V' \to \mathbb{A}^{m+n}$ satisfying the analogues of (a)–(d) in §4.3, where (d) becomes $\check{g} \circ \jmath' = \check{f} \circ \jmath' + z_1^2 + \cdots + z_n^2$.

Set $U' = \{v' \in V' : z_1(v') = \cdots = z_n(v') = 0\}$, which is a smooth $\mathbb{K}$-subscheme of V' as $(y_1,\ldots,y_m,z_1,\ldots,z_n)$ are étale coordinates on V'. Define $\iota : U' \to U$, $\jmath : V' \to V$, $\Phi' : U' \to V'$, $\alpha : V' \to U$, and $\beta : V' \to \mathbb{A}^n$ by $\iota = \check{\alpha} \circ \jmath'|_{U'}$, $\jmath = \check{\jmath} \circ \jmath'$, $\Phi' = \mathrm{id}_{U'}$, $\alpha = \check{\alpha} \circ \jmath'$ and $\beta = (z_1,\ldots,z_n)$. As $\jmath' : V' \to \check{V}$ is an étale open neighbourhood of $\check{v}$ in $\check{V}$, there exists $u' \in V'$ with $\jmath'(u') = \check{v}$, and $z_b(u') = 0$ for $b = 1,\ldots,n$ as $\check{z}_b(\check{v}) = \tilde{z}_b \circ \check{\jmath}(\check{v}) = \tilde{z}_b(j(x)) = 0$, so $u' \in U$ with $\iota(u') = \check{\alpha} \circ \jmath'(u') = \check{\alpha}(\check{v}) = i(x)$. Also $\iota, \jmath, \alpha \times \beta$ are étale as $\check{\alpha} \times \check{\beta}, \check{\jmath}, \jmath'$ are.

To see that $\Phi \circ \iota = \jmath \circ \Phi'$, note that

$$\begin{aligned}
&(\tilde{y}_1,\ldots,\tilde{y}_m,\tilde{z}_1,\ldots,\tilde{z}_n) \circ \Phi \circ \iota = (\tilde{x}_1,\ldots,\tilde{x}_m,0,\ldots,0) \circ \check{\alpha} \circ \jmath'|_{U'} \\
&= \big((\tilde{x}_1,\ldots,\tilde{x}_m) \times \mathrm{id}_{\mathbb{A}^n}\big) \circ (\check{\alpha} \times \check{\beta}) \circ \jmath'|_{U'} \\
&= (\tilde{y}_1,\ldots,\tilde{y}_m,\tilde{z}_1,\ldots,\tilde{z}_n) \circ \check{\jmath} \circ \jmath'|_{U'} \circ \mathrm{id}_{U'} = (\tilde{y}_1,\ldots,\tilde{y}_m,\tilde{z}_1,\ldots,\tilde{z}_n) \circ \jmath \circ \Phi',
\end{aligned}$$

using the definitions and (4.12). Since $\Phi\circ\iota(u')=j(x)=\jmath\circ\Phi'(u')$ and $(\tilde y_1,\ldots,\tilde y_m,\tilde z_1,\ldots,\tilde z_n)$ are étale coordinates on $\tilde V\subseteq V$, this implies that $\Phi\circ\iota=\jmath\circ\Phi'$ near u' in U', so making U' smaller if necessary we have $\Phi\circ\iota=\jmath\circ\Phi'$. The equations $\alpha\circ\Phi'=\iota$, $\beta\circ\Phi'=0$ are immediate, and $g\circ\jmath=f\circ\alpha+(z_1^2+\cdots+z_n^2)\circ\beta$ follows from $\check g\circ\jmath'=\check f\circ\jmath'+z_1^2+\cdots+z_n^2$. This proves Proposition 2.16.

4.5 Proof of Proposition 2.17

For Proposition 2.17, we first follow the proof of Proposition 2.16 in §4.4 until immediately before the choice of $\jmath':V'\to\check V$, so we have étale coordinates $(\tilde y_1,\ldots,\tilde y_m,\tilde z_1,\ldots,\tilde z_n)$ on $j(x)\in\tilde V\subseteq V$ and $(\tilde x_1,\ldots,\tilde x_m)$ on $i(x)\in\tilde U=\Phi^{-1}(\tilde V)\subseteq U$ with $\tilde x_a=\tilde y_a\circ\Phi|_{\tilde U}$, $\frac{\partial g}{\partial\tilde z_b}|_{\Phi(\tilde U)}=0$, and a Cartesian square (4.12) with étale coordinates $(\check y_1,\ldots,\check y_m,\check z_1,\ldots,\check z_n)$ on $\check V$ and $\check v\in\check V$ with $\check\alpha(\check v)=i(x)$, $\check\beta(\check v)=(0,\ldots,0)$ and $\check\jmath(\check v)=j(x)$, and functions $\check f,\check g,\check h,Q_{bc}:\check V\to\mathbb{A}^1$ with $\check f=f\circ\check\alpha$, $\check g=g\circ\check\jmath$, and $\check h=\check g-\check f=\sum_{b,c=1}^n\check z_b\check z_cQ_{bc}$, with $Q_{bc}(\check v)=\delta_{bc}$.

We have morphisms $\mathrm{id}\times 0:\tilde U\to\tilde U\times\mathbb{A}^n$, $\Phi|_{\tilde U}:\tilde U\to\tilde V$ with $\bigl((\tilde x_1,\ldots,\tilde x_m)\times\mathrm{id}_{\mathbb{A}^n}\bigr)\circ(\mathrm{id}\times 0)=(\tilde y_1,\ldots,\tilde y_m,\tilde z_1,\ldots,\tilde z_n)\circ\Phi|_{\tilde U}:\tilde U\to\mathbb{A}^{m+n}$. Thus by the Cartesian property of (4.12), there is a unique morphism $\check\Phi:\tilde U\to\check V$ with $(\check\alpha\times\check\beta)\circ\check\Phi=\mathrm{id}\times 0$ and $\check\jmath\circ\check\Phi=\Phi|_{\tilde U}$. Also $\check\Phi(i(x))=\check v$, since $(\mathrm{id}\times 0)(i(x))=(i(x),0)=(\check\alpha\times\check\beta)(\check v)$ and $\Phi|_{\tilde U}(i(x))=j(x)=\check\jmath(\check v)$.

We now modify the inductive procedure in §4.3–§4.4, to construct a Zariski open neighbourhood U' of $i(x)$ in $\tilde U\subseteq U$, an étale open neighbourhood $\jmath':V'\to\check V$ of $\check v$ in $\check V$ with $\check\alpha\circ\jmath'(V')\subseteq U'\subseteq\tilde U$, a morphism $\Phi':U'\to V'$ with $\jmath'\circ\Phi'=\check\Phi|_{U'}$, étale coordinates $(y_1,\ldots,y_m,z_1,\ldots,z_n):V'\to\mathbb{A}^{m+n}$, and regular functions $q_1,\ldots,q_n:U'\to\mathbb{A}^1\setminus\{0\}$, such that:

(a) $y_a=\check y_a\circ\jmath'=\tilde y_a\circ\check\jmath\circ\jmath'$ for $a=1,\ldots,m$.

(b) $\frac{\partial z_b}{\partial z'_c}=\delta_{bc}$ at $\Phi'(i(x))$ for $b,c=1,\ldots,n$, where $(y'_1,\ldots,y'_m,z'_1,\ldots,z'_n)=(\check y_1,\ldots,\check y_m,\check z_1,\ldots,\check z_n)\circ\jmath'$ are étale coordinates on V'.

(c) $\Phi'(U')$ is the submanifold $z_1=\cdots=z_n=0$ in V'.

(d) $\check h\circ\jmath'=(q_1\circ\check\alpha\circ\jmath')\cdot z_1^2+\cdots+(q_n\circ\check\alpha\circ\jmath')\cdot z_n^2$.

In the first step of the induction, as $Q_{nn}(\check v)=1$ and $\check\Phi(i(x))=\check v$, we can choose a Zariski open neighbourhood U' of $i(x)$ in $\tilde U$ such that $q_n:=Q_{nn}\circ\check\Phi$ is nonzero on U', so that $q_n:U'\to\mathbb{A}^1\setminus\{0\}$. Then $\check\alpha^{-1}(U')$ is a Zariski open neighbourhood of $\check v$ in $\check V$, with $\check\Phi(U')\subseteq\check\alpha^{-1}(U')$ as $\check\alpha\circ\check\Phi=\mathrm{id}_{\tilde U}$, and

$$\check P_n:=\bigl(Q_{nn}|_{\check\alpha^{-1}(U')}\bigr)\cdot\bigl(q_n^{-1}\circ\check\alpha|_{\check\alpha^{-1}(U')}\bigr):\check\alpha^{-1}(U')\longrightarrow\mathbb{A}^1$$

is a regular function, with $\check P_n\circ\check\Phi|_{U'}=1$.

Define $\hat\jmath:\hat V\to\check P_n^{-1}(\mathbb{A}^1\setminus\{0\})\subseteq\check V$ to be the étale double cover parametrizing square roots of $\check P_n$ wherever $\check P_n$ is nonzero. Then $\check P_n\circ\hat\jmath$ has a natural square root $(\check P_n\circ\hat\jmath)^{1/2}:\hat V\to\mathbb{A}^1\setminus\{0\}$. Since $\check P_n\circ\check\Phi|_{U'}=1$, there is a unique lift $\hat\Phi:U'\to\hat V$ such that $\hat\jmath\circ\hat\Phi=\check\Phi|_{U'}$ and $(\check P_n\circ\hat\jmath)^{1/2}\circ\hat\Phi=1$.

In an analogue of (4.11), we may now write

$$\begin{aligned}\check h\circ\hat\jmath&=\textstyle\sum_{b,c=1}^{n}(\check z_b\check z_cQ_{bc})\circ\hat\jmath=\sum_{b,c=1}^{n-1}\bigl(\check z_b\check z_c\bigl[Q_{bc}-Q_{nn}^{-1}Q_{bn}Q_{cn}\bigr]\bigr)\circ\hat\jmath\\&\quad+(q_n\circ\check\alpha\circ\hat\jmath)\cdot\Bigl[(\check P_n\circ\hat\jmath)^{1/2}\bigl(\check z_n+\textstyle\sum_{b=1}^{n-1}Q_{nn}^{-1}Q_{bn}\check z_b\bigr)\circ\hat\jmath\Bigr]^2\\&=\textstyle\sum_{b,c=1}^{n-1}\hat z_b\hat z_c\hat Q_{bc}+(q_n\circ\hat\alpha)\cdot\hat z_n^2,\end{aligned}\tag{4.13}$$

where $\hat z_b=\check z_b\circ\hat\jmath$ for $b=1,\ldots,n-1$, $\hat z_n=(\check P_n\circ\hat\jmath)^{1/2}\bigl(\check z_n+\sum_{b=1}^{n-1}Q_{nn}^{-1}Q_{bn}\check z_b\bigr)\circ\hat\jmath$, $\hat Q_{bc}=[Q_{bc}-Q_{nn}^{-1}Q_{bn}Q_{cn}]\circ\hat\jmath$ for $b,c=1,\ldots,n-1$, and $\hat\alpha=\check\alpha\circ\hat\jmath$.

In the second step we define $q_{n-1}:=\hat Q_{n-1\,n-1}\circ\hat\Phi$, and making U' smaller we can suppose that q_{n-1} maps $U'\to\mathbb{A}^1\setminus\{0\}$. Then we define

$$\hat P_{n-1}:=\bigl(\hat Q_{n-1\,n-1}|_{\hat\alpha^{-1}(U')}\bigr)\cdot\bigl(q_{n-1}^{-1}\circ\hat\alpha|_{\hat\alpha^{-1}(U')}\bigr):\hat\alpha^{-1}(U')\longrightarrow\mathbb{A}^1,$$

so that $\hat P_{n-1}\circ\hat\Phi=1$. Let $\ddot\jmath:\ddot V\to\hat P_{n-1}^{-1}(\mathbb{A}^1\setminus\{0\})\subseteq\hat V$ be the étale double cover parametrizing square roots of $\hat P_{n-1}$ where $\hat P_{n-1}$ is nonzero. Then $\hat P_{n-1}\circ\ddot\jmath$ has a square root $(\hat P_{n-1}\circ\ddot\jmath)^{1/2}:\ddot V\to\mathbb{A}^1\setminus\{0\}$, and there is a unique lift $\ddot\Phi:U'\to\ddot V$ such that $\ddot\jmath\circ\ddot\Phi=\hat\Phi$ and $(\hat P_{n-1}\circ\ddot\jmath)^{1/2}\circ\ddot\Phi=1$. As for (4.13) we have

$$\check h\circ\hat\jmath\circ\ddot\jmath=\textstyle\sum_{b,c=1}^{n-2}\ddot z_b\ddot z_c\ddot Q_{bc}+(q_{n-1}\circ\ddot\alpha)\cdot\ddot z_{n-1}^2+(q_n\circ\ddot\alpha)\cdot\ddot z_n^2,$$

where $\ddot z_{n-1}=(\hat P_{n-1}\circ\ddot\jmath)^{1/2}\bigl(\hat z_{n-1}+\sum_{b=1}^{n-2}\hat Q_{n-1n-1}^{-1}\hat Q_{bn-1}\hat z_b\bigr)\circ\ddot\jmath$, $\ddot z_b=\hat z_b\circ\ddot\jmath$ for $b=1,\ldots,n-2,n$, $\ddot Q_{bc}=[\hat Q_{bc}-\hat Q_{n-1n-1}^{-1}\hat Q_{bn-1}\hat Q_{cn-1}]\circ\ddot\jmath$ for $b,c=1,\ldots,n-2$, and $\ddot\alpha=\hat\alpha\circ\ddot\jmath$. Continuing in this way, after n inductive steps we define data $\jmath',V',\Phi',y_a,z_b,q_b$ satisfying (a)–(d) above. The important difference with §4.4 is that each time we pass to a further étale cover $\hat V,\ddot V,\ldots$ of $\check V$ to take a square root, we also lift $\check\Phi|_{U'}:U'\to\check V$ to morphisms $\hat\Phi:U'\to\hat V$, $\ddot\Phi:U'\to\ddot V$, …, to these étale covers, for $U'\subseteq U$ Zariski open. Proposition 2.17 now follows as in §4.3–§4.4, with (2.11) coming from (d) above.

5 Canonical bundles of d-critical loci

Finally we prove the results of §2.4, Propositions 2.18, 2.20 and Theorem 2.21.

5.1 Proof of Proposition 2.18

We will prove the complex analytic case of Proposition 2.18, involving (a) and Proposition 2.15. The algebraic cases with (b),(c) and Propositions 2.16 and 2.17 are similar.

Let (X,s) be a complex analytic d-critical locus and x,U',V',n,α,β be as in Proposition 2.15. To see there exists a unique isomorphism $\hat\beta$ satisfying (2.13), consider the diagram of vector bundles on U':

$$\begin{tikzcd}
0 \arrow[r] & TU|_{U'} \arrow[r, "\mathrm{d}\Phi|_{U'}"'] \arrow[d, "\cong"', "\mathrm{id}"] & \Phi^*(TV)|_{U'} \arrow[r, "\Pi|_{U'}"'] \arrow[d, "\cong"', "\Phi|_{U'}^*(\mathrm{d}(\alpha\times\beta))"] & N_{UV}|_{U'} \arrow[r] \arrow[d, dotted, "\cong"', "\hat\beta^*"] & 0 \\
0 \arrow[r] & TU|_{U'} \arrow[r, "\mathrm{d}(\mathrm{id}\times 0)"] & T(U\times\mathbb{C}^n)|_{U'\times 0} \arrow[r] & \bigl\langle\frac{\partial}{\partial z_1},\ldots,\frac{\partial}{\partial z_n}\bigr\rangle_{U'} \arrow[r] & 0.
\end{tikzcd}\tag{5.1}$$

Here (5.1) has exact rows, the first two columns are isomorphisms as $\alpha \times \beta$ is étale, and the left hand square commutes since $\alpha \circ \Phi|_{U'} = \mathrm{id}$ and $\beta \circ \Phi|_{U'} = 0$. Therefore by exactness there is a unique isomorphism $\hat{\beta}^*$ making (5.1) commute, and taking duals shows that (2.13) holds. So (2.14) prescribes $q_{UV}|_{R'}$, for $R' = i^{-1}(U') \subseteq R \subseteq X$. This $q_{UV}|_{R'}$ is a nondegenerate holomorphic quadratic form on $i^*(N_{UV})|_{R'}$, since $\mathrm{d}z_1 \otimes \mathrm{d}z_1 + \cdots + \mathrm{d}z_n \otimes \mathrm{d}z_n$ is a nondegenerate holomorphic quadratic form on $\langle \frac{\partial}{\partial z_1}, \ldots, \frac{\partial}{\partial z_n} \rangle_{U'}$.

Thus, for each $x \in R$ we can find an open $x \in R' \subseteq R$ and a given value for the restriction $q_{UV}|_{R'}$. Since such R' form an open cover for R, and $S^2 i^*(N_{UV}^*)$ is a sheaf on R, these values for $q_{UV}|_{R'}$ come from a unique $q_{UV} \in H^0(S^2 i^*(N_{UV}^*))$ if and only if they agree on overlaps $R' \cap R''$ between different subsets R', R'' in the open cover.

Suppose $x, U', V', \alpha, \beta, R'$ and $x', U'', V'', \alpha', \beta', R''$ are alternative choices in Proposition 2.15. Then (2.14) gives values for $q_{UV}|_{R'}$ and $q_{UV}|_{R''}$, which agree on the overlap $R' \cap R''$ if

$$\begin{aligned} &i|_{R'\cap R''}^* \big[(S^2\hat{\beta})(\mathrm{d}z_1 \otimes \mathrm{d}z_1 + \cdots + \mathrm{d}z_n \otimes \mathrm{d}z_n)\big] \\ &\quad = i|_{R'\cap R''}^* \big[(S^2\hat{\beta}')(\mathrm{d}z_1 \otimes \mathrm{d}z_1 + \cdots + \mathrm{d}z_n \otimes \mathrm{d}z_n)\big]. \end{aligned} \tag{5.2}$$

Combining (5.1) for U', V', α, β and $U'', V'', \alpha', \beta'$ gives a commutative diagram of vector bundles on $U' \cap U''$, with exact rows:

$$\begin{tikzcd}
0 \arrow[r] & TU|_{U'} \arrow[r, "\mathrm{d}(\mathrm{id}\times 0)"'] \arrow[d, "\mathrm{id}", "\cong"'] & T(U\times\mathbb{C}^n)|_{U'\times 0} \arrow[r, "\Pi|_{U'}"'] \arrow[d, "{\Phi|_{U'}^*(\mathrm{d}((\alpha'\times\beta') \circ(\alpha\times\beta)^{-1}))}", "\cong"'] & \big\langle \frac{\partial}{\partial z_1}, \ldots, \frac{\partial}{\partial z_n} \big\rangle_{U'} \arrow[r] \arrow[d, "\hat{\beta}'^*\circ\hat{\beta}^{*-1}", "\cong"'] & 0 \\
0 \arrow[r] & TU|_{U'} \arrow[r, "\mathrm{d}(\mathrm{id}\times 0)"] & T(U\times\mathbb{C}^n)|_{U'\times 0} \arrow[r] & \big\langle \frac{\partial}{\partial z_1}, \ldots, \frac{\partial}{\partial z_n} \big\rangle_{U'} \arrow[r] & 0.
\end{tikzcd} \tag{5.3}$$

Here $(\alpha' \times \beta') \circ (\alpha \times \beta)^{-1}$ is a local biholomorphism $U \times \mathbb{C}^n \to U \times \mathbb{C}^n$ defined near $(U' \cap U'') \times \{0\}$, which is the identity on $U \times \{0\}$, and preserves the function $f \boxplus z_1^2 + \cdots + z_n^2 : U \times \mathbb{C}^n \to \mathbb{C}$. So restricting to $\mathrm{Crit}(f \boxplus z_1^2 + \cdots + z_n^2) = \mathrm{Crit}(f) \times \{0\}$ where $\mathrm{Hess}\big(f \boxplus z_1^2 + \cdots + z_n^2\big)$ is defined, we see that $\mathrm{d}((\alpha' \times \beta') \circ (\alpha \times \beta)^{-1})|_{\mathrm{Crit}(f|_{U'\cap U''})\times\{0\}}$ preserves $\mathrm{Hess}\big(f \boxplus z_1^2 + \cdots + z_n^2\big)$ in $H^0\big(S^2 T^*(U \times \mathbb{C}^n)|_{\mathrm{Crit}(f|_{U'\cap U''})\times\{0\}}\big)$. As

$$\mathrm{Hess}\big(f \boxplus z_1^2 + \cdots + z_n^2\big) = \mathrm{Hess}\, f + \mathrm{d}z_1 \otimes \mathrm{d}z_1 + \cdots + \mathrm{d}z_n \otimes \mathrm{d}z_n,$$

from (5.3) we see that $\hat{\beta}'^* \circ \hat{\beta}^{*-1}$ preserves $\mathrm{d}z_1 \otimes \mathrm{d}z_1 + \cdots + \mathrm{d}z_n \otimes \mathrm{d}z_n$ on $\mathrm{Crit}(f|_{U'\cap U''})$. That is,

$$\begin{aligned} &(S^2(\hat{\beta}^{-1} \circ \hat{\beta}'))(\mathrm{d}z_1 \otimes \mathrm{d}z_1 + \cdots + \mathrm{d}z_n \otimes \mathrm{d}z_n)|_{\mathrm{Crit}(f|_{U'\cap U''})} \\ &\qquad = (\mathrm{d}z_1 \otimes \mathrm{d}z_1 + \cdots + \mathrm{d}z_n \otimes \mathrm{d}z_n)|_{\mathrm{Crit}(f|_{U'\cap U''})}. \end{aligned}$$

Composing with $S^2\hat{\beta}$ and applying $i|_{R'\cap R''}^*$ gives (5.2). Hence, there exists a unique, nondegenerate q_{UV} satisfying Proposition 2.18(a).

For the final part of the proposition, suppose $\Psi : (S, V, g, j) \hookrightarrow (T, W, h, k)$ is another embedding, so that $\Psi \circ \Phi : (R, U, f, i) \hookrightarrow (T, W, h, k)$ is also an embedding, and define N_{UV}, q_{UV}, N_{VW}, q_{VW} and N_{UW}, q_{UW} from Φ, Ψ and $\Psi \circ \Phi$ as above. Then existence of unique $\gamma_{UVW}, \delta_{UVW}$ making (2.19) commute is immediate by exactness, and exactness of the line in (2.19) including $\gamma_{UVW}, \delta_{UVW}$ also follows easily. So (2.20) is exact on R.

We will first prove the required isomorphism (2.21) exists locally. Let $x \in R$. Applying Proposition 2.15 to $\Phi : (R, U, f, i) \hookrightarrow (S, V, g, j)$ at x gives $i(x) \in U' \subseteq U$, $j(x) \in V' \subseteq V$, $\alpha : V' \to U$ and $\beta : V' \to \mathbb{C}^m$ for $m = \dim V - \dim U$, satisfying conditions including $g|_{V'} = f \circ \alpha + (y_1^2 + \cdots + y_m^2) \circ \beta$, writing $(y_1, \ldots, y_m)$ for the coordinates on $\mathbb{C}^m$. Then Proposition 2.18(a) gives

$$\begin{gathered}\hat\beta : \langle \mathrm{d}y_1, \ldots, \mathrm{d}y_m \rangle_{U'} \xrightarrow{\cong} N_{UV}^*|_{U'}, \qquad \text{and} \\ q_{UV}|_{R'} = i|_{R'}^* \bigl[(S^2\hat\beta)(\mathrm{d}y_1 \otimes \mathrm{d}y_1 + \cdots + \mathrm{d}y_m \otimes \mathrm{d}y_m)\bigr],\end{gathered} \tag{5.4}$$

where $R' = i^{-1}(U') \subseteq R \subseteq X$. Similarly, applying Proposition 2.15 to $\Psi : (S, V, g, j) \hookrightarrow (T, W, h, k)$ at x gives $j(x) \in V'' \subseteq V$, $k(x) \in W'' \subseteq V$, $\alpha' : W'' \to V$ and $\beta' : W'' \to \mathbb{C}^n$ for $n = \dim W - \dim V$, satisfying conditions including $h|_{W''} = g \circ \alpha' + (z_1^2 + \cdots + z_n^2) \circ \beta'$, writing $(z_1, \ldots, z_n)$ for the coordinates on $\mathbb{C}^n$, and with $S'' = j^{-1}(V'')$, Proposition 2.18(a) gives

$$\begin{gathered}\hat\beta' : \langle \mathrm{d}z_1, \ldots, \mathrm{d}z_n \rangle_{V''} \xrightarrow{\cong} N_{VW}^*|_{V''}, \qquad \text{and} \\ q_{VW}|_{S''} = j|_{S''}^* \bigl[(S^2\hat\beta')(\mathrm{d}z_1 \otimes \mathrm{d}z_1 + \cdots + \mathrm{d}z_n \otimes \mathrm{d}z_n)\bigr].\end{gathered} \tag{5.5}$$

Set $U''' = U' \cap \Phi^{-1}(V'')$ and $W''' = \alpha'^{-1}(V')$, write $(y_1, \ldots, y_m, z_1, \ldots, z_n)$ for the coordinates on $\mathbb{C}^{m+n}$, and define $\alpha'' : W''' \to U$ and $\beta'' : W''' \to \mathbb{C}^{m+n}$ by $\alpha'' = \alpha \circ \alpha'|_{W'''}$ and $\beta'' = (\beta \circ \alpha'|_{W'''}) \times \beta'|_{W'''}$. Then

$$\begin{aligned} h|_{W'''} &= g \circ \alpha'|_{W'''} + (z_1^2 + \cdots + z_n^2) \circ \beta'|_{W'''} \\ &= f \circ \alpha \circ \alpha'|_{W'''} + (y_1^2 + \cdots + y_m^2) \circ \beta \circ \alpha'|_{W'''} + (z_1^2 + \cdots + z_n^2) \circ \beta'|_{W'''} \\ &= f \circ \alpha'' + (y_1^2 + \cdots + y_m^2 + z_1^2 + \cdots + z_n^2) \circ \beta''. \end{aligned}$$

The other conditions are easy to verify, so $U''', W''', \alpha'', \beta'', m+n$ are a possible outcome for Proposition 2.15 applied to $\Psi \circ \Phi : (R, U, f, i) \hookrightarrow (T, W, h, k)$ at x. Hence Proposition 2.18(a) with $R''' = i^{-1}(U''') = R' \cap S''$ gives

$$\begin{aligned} &\hat\beta'' : \langle \mathrm{d}y_1, \ldots, \mathrm{d}y_m, \mathrm{d}z_1, \ldots, \mathrm{d}z_n \rangle_{W'''} \xrightarrow{\cong} N_{UW}^*|_{W'''}, \qquad \text{and} \\ &q_{UW}|_{R'''} = i|_{W'''}^* \bigl[(S^2\hat\beta'')(\mathrm{d}y_1 \otimes \mathrm{d}y_1 + \cdots + \mathrm{d}y_m \otimes \mathrm{d}y_m \\ &\qquad\qquad + \mathrm{d}z_1 \otimes \mathrm{d}z_1 + \cdots + \mathrm{d}z_n \otimes \mathrm{d}z_n)\bigr]. \end{aligned} \tag{5.6}$$

The isomorphism $i^*(N_{UW}) \cong i^*(N_{UV}) \oplus j^*(N_{VW})|_R$ in (2.21) is now clear on the open subset $x \in R''' = R' \cap S'' \subseteq R$, from the isomorphisms

$$\begin{aligned} i|_{R'''}^*(\hat\beta) &: \langle \mathrm{d}y_1, \ldots, \mathrm{d}y_m \rangle_{R'''} \xrightarrow{\cong} i^*(N_{UV})|_{R'''}^*, \\ j|_{R'''}^*(\hat\beta') &: \langle \mathrm{d}z_1, \ldots, \mathrm{d}z_n \rangle_{R'''} \xrightarrow{\cong} j^*(N_{VW})|_{R'''}^*, \\ i|_{R'''}^*(\hat\beta'') &: \langle \mathrm{d}y_1, \ldots, \mathrm{d}y_m \rangle_{R'''} \oplus \langle \mathrm{d}z_1, \ldots, \mathrm{d}z_n \rangle_{R'''} \xrightarrow{\cong} i^*(N_{UW})|_{R'''}^*. \end{aligned}$$

It is easy to see this isomorphism is compatible with (2.20), and (5.4)–(5.6) imply that the isomorphism induces equation (2.22) on R'''.

Now isomorphisms (2.21) compatible with (2.20) are in 1-1 correspondence with complementary vector subbundles to $i^*(\gamma_{UVW})\bigl(i^*(N_{UV})\bigr)$ in $i^*(N_{UW})$. In this case, the complementary vector subbundle is just the orthogonal subbundle to $i^*(\gamma_{UVW})\bigl(i^*(N_{UV})\bigr)$ using the complex inner product q_{UW} on $i^*(N_{UW})$. This orthogonal subbundle is complementary provided $q_{UW}|_{i^*(\gamma_{UVW})(i^*(N_{UV}))}$ is nondegenerate, which holds as this restriction is isomorphic to q_{UV}.

Thus, the isomorphism (2.21) exists locally on R, and is unique (even locally) if it exists. So we can glue local choices on an open cover of R by subsets R''' to get a unique global isomorphism (2.21) compatible with (2.20) and (2.22). This completes the proof of Proposition 2.18.

5.2 Proof of Proposition 2.20

Let $\Phi : (R,U,f,i) \hookrightarrow (S,V,g,j)$ be an embedding of critical charts on a complex analytic d-critical locus (X,s). Fix $x \in R \subseteq X$. As for (2.12), define the normal $\mathbb{C}$-vector spaces $N_{XU}|_x, N_{XV}|_x$ to X in U,V at x by the exact sequences

$$\begin{tikzcd}
0 \arrow[r] & T_xX \arrow[r, "\mathrm{d}i|_x"'] & T_{i(x)}U \arrow[r] & N_{XU}|_x \arrow[r] & 0, \\
0 \arrow[r] & T_xX \arrow[r, "\mathrm{d}j|_x"] & T_{j(x)V} \arrow[r] & N_{XV}|_x \arrow[r] & 0,
\end{tikzcd} \tag{5.7}$$

where T_xX is the Zariski tangent space of X at x. Write $\dim T_xX = l$, $\dim U = l+m$ and $\dim V = l+m+n$, so that $\dim N_{XU}|_x = m$, $\dim N_{XV}|_x = m+n$, and $\dim N_{UV}|_{i(x)} = n$. As for (2.23), equation (5.7) induces isomorphisms

$$\begin{aligned}
&\rho_{XU}|_x : \Lambda^l T_x^*X \otimes \Lambda^m N_{XU}^*|_x \longrightarrow \Lambda^{l+m} T_{i(x)}^*U = K_U|_{i(x)}, \\
&\rho_{XV}|_x : \Lambda^l T_x^*X \otimes \Lambda^{m+n} N_{XV}^*|_x \longrightarrow \Lambda^{l+m+n} T_{j(x)}^*V = K_V|_{j(x)}.
\end{aligned} \tag{5.8}$$

We also have a commutative diagram with exact rows:

$$\begin{tikzcd}
0 \arrow[r] & T_xX \arrow[r, "\mathrm{d}i|_x"'] \arrow[d, "\mathrm{id}"'] & T_{i(x)}U \arrow[r, "\mathrm{Hess}_{i(x)} f"'] \arrow[d, "\mathrm{d}\Phi|_{i(x)}"] & T_{i(x)}^*U \arrow[r, "\mathrm{d}i|_x^*"'] \arrow[d, dotted] & T_x^*X \arrow[r] \arrow[d, "\mathrm{id}"] & 0 \\
0 \arrow[r] & T_xX \arrow[r, "\mathrm{d}j|_x"] & T_{j(x)}V \arrow[r, "\mathrm{Hess}_{j(x)} g"] & T_{j(x)}^*V \arrow[r, "\mathrm{d}j|_x^*"] & T_x^*X \arrow[r] & 0.
\end{tikzcd} \tag{5.9}$$

Since $T_xX = \mathrm{Ker}(\mathrm{Hess}_{i(x)} f)$, by (5.7) $\mathrm{Hess}_{i(x)} f$ is the pullback to $T_{i(x)}U$ of a nondegenerate quadratic form $\mathrm{Hess}'_{i(x)} f$ on $N_{XU}|_x$. Then $\det(\mathrm{Hess}'_{i(x)} f)$ is a nonzero element of $\Lambda^m N_{XU}^*|_x^{\otimes^2}$. Similarly, $\mathrm{Hess}_{j(x)} g$ is the pullback to $T_{j(x)}V$ of $\mathrm{Hess}'_{j(x)} g$ on $N_{XV}|_x$, and $0 \neq \det(\mathrm{Hess}'_{j(x)} g) \in \Lambda^{m+n} N_{XV}^*|_x^{\otimes^2}$.

In a similar way to (2.20), there is a natural exact sequence

$$\begin{tikzcd}
0 \arrow[r] & N_{XU}|_x \arrow[r] & N_{XV}|_x \arrow[r] & N_{UV}|_{i(x)} \arrow[r] & 0,
\end{tikzcd} \tag{5.10}$$

and as for (2.21), there is a unique isomorphism $N_{XV}|_x \cong N_{XU}|_x \oplus N_{UV}|_{i(x)}$ compatible with (5.10) and identifying $\mathrm{Hess}'_{j(x)}\, g$ with $\mathrm{Hess}'_{i(x)}\, f \oplus q_{UV}|_x \oplus 0$ as in (2.22). From this, we deduce that the following diagram commutes:

$$\begin{tikzcd}
(\Lambda^{l+m}T^*_{i(x)}U)^{\otimes^2} = K_U^{\otimes^2}|_{i(x)} \arrow[r, "\mathrm{id}\otimes\det(q_{UV})|_x"'] \arrow[dr, "J_\Phi|_x"'] & K_U^{\otimes^2}|_{i(x)} \otimes \Lambda^n N^*_{UV}|^{\otimes^2}_{i(x)} \arrow[d, "\rho^{\otimes^2}_{UV}|_x"'] \\
(\Lambda^l T^*_x X)^{\otimes^2} \otimes \Lambda^m N^*_{XU}|^{\otimes^2}_x \arrow[u, "\rho^{\otimes^2}_{XU}|_x"] & (\Lambda^{l+m+n}T^*_{j(x)}V)^{\otimes^2} = K_V^{\otimes^2}|_{j(x)} \\
(\Lambda^l T^*_x X)^{\otimes^2} \arrow[u, "\mathrm{id}\otimes\det(\mathrm{Hess}'_{i(x)}\, f)"] \arrow[r, "\mathrm{id}\otimes\det(\mathrm{Hess}'_{j(x)}\, g)"] & (\Lambda^l T^*_x X)^{\otimes^2} \otimes \Lambda^{m+n} N^*_{XV}|^{\otimes^2}_x . \arrow[u, "\rho^{\otimes^2}_{XV}|_x"']
\end{tikzcd} \tag{5.11}$$

Here the upper right triangle is the restriction of (2.24) to x, and depends on Φ. But the rest of the diagram depends on $x, (R, U, f, i), (S, V, g, j)$ but not on Φ. So (5.11) a diagram of commuting isomorphisms implies that $J_\Phi|_x$ is independent of the choice of Φ. Thus, if $\Phi, \tilde\Phi : (R, U, f, i) \hookrightarrow (S, V, g, j)$ are embeddings of critical charts, then $J_\Phi|_x = J_{\tilde\Phi}|_x$ for all $x \in R^{\mathrm{red}}$. As R^{red} is a reduced complex analytic space, this implies that $J_\Phi = J_{\tilde\Phi}$, as we want.

To prove (2.25), consider the commutative diagram

$$\begin{tikzcd}
 & & i^*(K_U^{\otimes^2})\otimes i^*(\Lambda^{\mathrm{top}}N^*_{UW})^{\otimes^2}|_{R^{\mathrm{red}}} \arrow[d, "\text{from (2.20)}"'] \arrow[ddd, bend left, "\rho^{\otimes^2}_{UW}"] \\
i^*(K_U^{\otimes^2})|_{R^{\mathrm{red}}} \arrow[urr, bend left=15, "\mathrm{id}\otimes\det(q_{UW})"] \arrow[r, "\mathrm{id}\otimes\det(q_{UV})"] \arrow[dr, "J_\Phi"'] \arrow[ddrr, bend right, "J_{\Psi\circ\Phi}"'] & i^*(K_U^{\otimes^2})\otimes i^*(\Lambda^{\mathrm{top}}N^*_{UV})^{\otimes^2}|_{R^{\mathrm{red}}} \arrow[r, "\mathrm{id}\otimes\det(q_{VW})"] \arrow[d, "\rho^{\otimes^2}_{UV}"] & u^*(K_U^{\otimes^2})\otimes i^*(\Lambda^{\mathrm{top}}N^*_{UV})^{\otimes^2}\otimes i^*(\Lambda^{\mathrm{top}}N^*_{VW})^{\otimes^2}|_{R^{\mathrm{red}}} \arrow[d, "\rho^{\otimes^2}_{UV}\otimes\mathrm{id}"'] \\
 & i^*(K_V^{\otimes^2})|_{R^{\mathrm{red}}} \arrow[r, "\mathrm{id}\otimes\det(q_{VW})"] \arrow[dr, "J_\Psi|_{R^{\mathrm{red}}}"'] & i^*(K_V^{\otimes^2})\otimes i^*(\Lambda^{\mathrm{top}}N^*_{VW})^{\otimes^2}|_{R^{\mathrm{red}}} \arrow[d, "\rho^{\otimes^2}_{VW}"'] \\
 & & k^*(K_W^{\otimes^2})|_{R^{\mathrm{red}}},
\end{tikzcd}$$

which includes (2.24) for Φ, Ψ and $\Psi \circ \Phi$, where the top left triangle commutes by (2.22). This proves Proposition 2.20 for complex analytic d-critical loci.

In the algebraic case, for (X, s) an algebraic d-critical locus over a field $\mathbb{K}$, the argument above shows that for each scheme-theoretic point x of the reduced $\mathbb{K}$-subscheme R^{red} we have $J_\Phi|_x = J_{\tilde\Phi}|_x$, where (5.7)–(5.11) are now diagrams of finite-dimensional vector spaces over the residue field of X at x, rather than over $\mathbb{C}$. Since R^{red} is a reduced $\mathbb{K}$-scheme and $J_\Phi, J_{\tilde\Phi}$ are isomorphisms of line bundles on R^{red}, $J_\Phi|_x = J_{\tilde\Phi}|_x$ for each $x \in R^{\mathrm{red}}$ implies that $J_\Phi = J_{\tilde\Phi}$. The rest of the proof is as for the complex analytic case.

5.3 Proof of Theorem 2.21

Observe that the structure of the first part of Theorem 2.1 characterizing the sheaf $\mathcal{S}_X$ by parts (i),(ii) and proved in §3.1, is very similar to the structure of Theorem 2.21 characterizing the sheaf $K_{X,s}$ by (i),(ii). We will follow the method of §3.1 to prove Theorem 2.21. We handle the complex analytic and algebraic cases together. The analogues of Lemmas 3.1, 3.2 and 3.3 are Proposition 2.20, Theorem 2.13 and:

Lemma 5.1. *Let $(R,U,f,i),(S,V,g,j)$ be critical charts on (X,s). Then there exists a unique isomorphism*

$$J_{R,U,f,i}^{S,V,g,j} : i^*\bigl(K_U^{\otimes^2}\bigr)|_{R^{\mathrm{red}}\cap S^{\mathrm{red}}} \longrightarrow j^*\bigl(K_V^{\otimes^2}\bigr)|_{R^{\mathrm{red}}\cap S^{\mathrm{red}}} \tag{5.12}$$

such that if $x,U'\subseteq U,V'\subseteq V,(T,W,h,k)$ and $\Phi:(R',U',f',i')\hookrightarrow(T,W,h,k)$, $\Psi:(S',V',g',j')\hookrightarrow(T,W,h,k)$ are as in Theorem 2.13, and J_Φ,J_Ψ are as in Definition 2.19 for Φ,Ψ, then

$$J_{R,U,f,i}^{S,V,g,j}|_{R'^{\mathrm{red}}\cap S'^{\mathrm{red}}} = J_\Psi^{-1}\circ J_\Phi|_{R'^{\mathrm{red}}\cap S'^{\mathrm{red}}}. \tag{5.13}$$

Also, if (T,W,h,k) is any other critical chart on (X,s) then

$$\begin{aligned} &J_{R,U,f,i}^{R,U,f,i} = \mathrm{id}_{i^*(K_U^2)|_{R^{\mathrm{red}}}}, \qquad J_{S,V,g,j}^{R,U,f,i} = \bigl(J_{R,U,f,i}^{S,V,g,j}\bigr)^{-1}, \\ \textit{and}\qquad &J_{S,V,g,j}^{T,W,h,k}\circ J_{R,U,f,i}^{S,V,g,j}|_{R^{\mathrm{red}}\cap S^{\mathrm{red}}\cap T^{\mathrm{red}}} = J_{R,U,f,i}^{T,W,h,k}|_{R^{\mathrm{red}}\cap S^{\mathrm{red}}\cap T^{\mathrm{red}}}. \end{aligned} \tag{5.14}$$

Proof. Suppose x, U', V', Φ, Ψ, (T,W,h,k) and $\check x$, $\check U'$, $\check V'$, $\check\Phi$, $\check\Psi$, $(\check T,\check W,\check h,\check k)$ are two possible choices in Theorem 2.13. We will show that

$$J_\Psi^{-1}\circ J_\Phi|_{R'^{\mathrm{red}}\cap S'^{\mathrm{red}}\cap\check R'^{\mathrm{red}}\cap\check S'^{\mathrm{red}}} = J_{\check\Psi}^{-1}\circ J_{\check\Phi}|_{R'^{\mathrm{red}}\cap S'^{\mathrm{red}}\cap\check R'^{\mathrm{red}}\cap\check S'^{\mathrm{red}}}. \tag{5.15}$$

Let $\tilde x\in R'^{\mathrm{red}}\cap S'^{\mathrm{red}}\cap\check R'^{\mathrm{red}}\cap\check S'^{\mathrm{red}}\subseteq T\cap\check T$. Then applying Theorem 2.13 to $\tilde x,(T,W,h,k),(\check T,\check W,\check h,\check k)$ yields open $\tilde x\in T'\subseteq T$, $\tilde x\in\check T'\subseteq\check T$, a critical chart $(\tilde T,\tilde W,\tilde h,\tilde k)$ and embeddings $\Theta:(T',W',h',k')\hookrightarrow(\tilde T,\tilde W,\tilde h,\tilde k)$ and $\Xi:(\check T',\check W',\check h',\check k')\hookrightarrow(\tilde T,\tilde W,\tilde h,\tilde k)$.

Set $U''=\Phi^{-1}(W')\cap\check\Phi^{-1}(\check W')$, $V''=\Psi^{-1}(W')\cap\check\Psi^{-1}(\check W')$, and let $(R'',U'',f'',i'')\subseteq(R,U,f,i)$, $(S'',V'',g'',j'')\subseteq(S,V,g,j)$ be the corresponding subcharts. Then we have a diagram of embeddings of critical charts

$$\begin{tikzcd}
(R'',U'',f'',i'') \arrow[r, "\Phi|_{U''}"'] \arrow[ddr, "\check\Phi|_{U''}"' near start] & (T',W',h',k') \arrow[dr, "\Theta"] & \\
 & & (\tilde T,\tilde W,\tilde h,\tilde k). \\
(S'',V'',g'',j'') \arrow[r, "\check\Psi|_{V''}"] \arrow[uur, "\Psi|_{V''}" near end] & (\check T',\check W',\check h',\check k') \arrow[ur, "\Xi"'] &
\end{tikzcd}$$

Hence we have

$$\begin{aligned} J_\Theta\circ J_\Phi|_{R''^{\mathrm{red}}} &= J_{\Theta\circ\Phi}|_{R''^{\mathrm{red}}} = J_{\Xi\circ\check\Phi}|_{R''^{\mathrm{red}}} = J_\Xi\circ J_{\check\Phi}|_{R''^{\mathrm{red}}}, \\ J_\Theta\circ J_\Psi|_{S''^{\mathrm{red}}} &= J_{\Theta\circ\Psi}|_{S''^{\mathrm{red}}} = J_{\Xi\circ\check\Psi}|_{S''^{\mathrm{red}}} = J_\Xi\circ J_{\check\Psi}|_{S''^{\mathrm{red}}}, \end{aligned} \tag{5.16}$$

using (2.25) in the first and third steps of each line, and the first part of Proposition 2.20 in the second steps. Restricting (5.16) to $R''^{\mathrm{red}} \cap S''^{\mathrm{red}}$, inverting the second line, and composing with the first line gives

$$\begin{aligned} J_\Psi^{-1} \circ J_\Phi|_{R''^{\mathrm{red}} \cap S''^{\mathrm{red}}} &= (J_\Theta \circ J_\Psi)^{-1} \circ (J_\Theta \circ J_\Phi)|_{R''^{\mathrm{red}} \cap S''^{\mathrm{red}}} \\ &= (J_\Xi \circ J_{\check\Psi})^{-1} \circ (J_\Xi \circ J_{\check\Phi})|_{R''^{\mathrm{red}} \cap S''^{\mathrm{red}}} = J_{\check\Psi}^{-1} \circ J_{\check\Phi}|_{R''^{\mathrm{red}} \cap S''^{\mathrm{red}}}. \end{aligned}$$

This proves (5.15) on the open neighbourhood $R''^{\mathrm{red}} \cap S''^{\mathrm{red}}$ of $\tilde x$ in $R'^{\mathrm{red}} \cap S'^{\mathrm{red}} \cap \check R'^{\mathrm{red}} \cap \check S'^{\mathrm{red}}$. As this works for all such $\tilde x$, equation (5.15) holds.

Thus, Theorem 2.13 shows that for each $x \in R^{\mathrm{red}} \cap S^{\mathrm{red}}$ we can choose an open neighbourhood $R'^{\mathrm{red}} \cap S'^{\mathrm{red}}$ on which the restriction of $J_{R,U,f,i}^{S,V,g,j}$ is defined by (5.13). These open neighbourhoods $R'^{\mathrm{red}} \cap S'^{\mathrm{red}}$ form an open cover of $R^{\mathrm{red}} \cap S^{\mathrm{red}}$, and on overlaps $(R'^{\mathrm{red}} \cap S'^{\mathrm{red}}) \cap (\check R'^{\mathrm{red}} \cap \check S'^{\mathrm{red}})$ the corresponding values of $J_{R,U,f,i}^{S,V,g,j}$ agree. Therefore by properties of sheaves there is a unique morphism $J_{R,U,f,i}^{S,V,g,j}$ in (5.12) such that (5.13) holds for all applications of Theorem 2.13. Finally, we prove (5.14) by the method used for (3.6)–(3.7) in Lemma 3.3. $\square$

Theorem 2.21 now follows from Lemma 5.1 in the same way that the first part of Theorem 2.1 was deduced from Lemma 3.3 in §3.1.

References

[1] K. Behrend, *Donaldson–Thomas type invariants via microlocal geometry*, Ann. of Math. 170 (2009), 1307–1338. math.AG/0507523.

[2] K. Behrend and B. Fantechi, *The intrinsic normal cone*, Invent. Math. 128 (1997), 45–88. alg-geom/9601010.

[3] C. Brav, V. Bussi, D. Dupont, D. Joyce, and B. Szendrői, *Symmetries and stabilization for sheaves of vanishing cycles*, arXiv:1211.3259v2, 2013.

[4] C. Brav, V. Bussi and D. Joyce, *A Darboux theorem for derived schemes with shifted symplectic structure*, arXiv:1305.6302, 2013.

[5] V. Bussi, D. Joyce and S. Meinhardt, *On motivic vanishing cycles of critical loci*, 1305.6428, 2013.

[6] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian intersection Floer theory — anomaly and obstruction*, Parts I & II. AMS/IP Studies in Advanced Mathematics, 46.1 & 46.2, A.M.S./International Press, 2009.

[7] H. Grauert and R. Remmert, *Coherent analytic sheaves*, Grundlehren der math. Wiss. 265, Springer-Verlag, Berlin, 1984.

[8] R.C. Gunning and H. Rossi, *Analytic functions of several complex variables*, Prentice-Hall, Englewood Cliffs, NJ, 1965.

[9] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Math. 52, Springer, New York, 1977.

[10] L. Illusie, *Complexe cotangent et déformations. I*, Springer Lecture Notes in Math. 239, Springer-Verlag, Berlin, 1971.

[11] L. Illusie, *Cotangent complex and deformations of torsors and group schemes*, pages 159–189 in Springer Lecture Notes in Math. 274, Springer-Verlag, Berlin, 1972.

[12] D. Joyce, *A classical model for derived Lagrangians in complex symplectic manifolds*, in preparation, 2013.

[13] D. Joyce and Y. Song, *A theory of generalized Donaldson–Thomas invariants*, Mem. Amer. Math. Soc. 217 (2012), no. 1020. arXiv:0810.5645.

[14] M. Kontsevich and Y. Soibelman, *Stability structures, motivic Donaldson–Thomas invariants and cluster transformations*, arXiv:0811.2435, 2008.

[15] M. Kontsevich and Y. Soibelman, *Cohomological Hall algebra, exponential Hodge structures and motivic Donaldson–Thomas invariants*, Commun. Number Theory Phys. 5 (2011), 231–352. arXiv:1006.2706.

[16] R. Pandharipande and R.P. Thomas, *Almost closed* 1*-forms*, arXiv:1204.3958, 2012.

[17] T. Pantev, B. Toën, M. Vaquié and G. Vezzosi, *Quantization and derived moduli spaces I: Shifted symplectic structures*, arXiv:1111.3209, 2011.

[18] J.-P. Serre, *Géométrie algébrique et géométrie analytique*, Ann. Inst. Fourier 6 (1956), 1–42.

[19] R.P. Thomas, *A holomorphic Casson invariant for Calabi–Yau* 3*-folds, and bundles on $K3$ fibrations*, J. Diff. Geom. 54 (2000), 367–438. math.AG/9806111.

[20] B. Toën, *Higher and derived stacks: a global overview*, pages 435–487 in *Algebraic Geometry — Seattle 2005*, Proc. Symp. Pure Math. 80, Part 1, A.M.S., Providence, RI, 2009. math.AG/0604504.

[21] B. Toën and G. Vezzosi, *Homotopical Algebraic Geometry II: Geometric Stacks and Applications*, Mem. Amer. Math. Soc. 193 (2008), no. 902. math.AG/0404373.

[22] B. Toën and G. Vezzosi, *From HAG to DAG: derived moduli stacks*, pages 173–216 in *Axiomatic, enriched and motivic homotopy theory*, NATO Sci. Ser. II Math. Phys. Chem., 131, Kluwer, Dordrecht, 2004. math.AG/0210407.

[23] G. Vezzosi, *Derived critical loci I: Basics*, arXiv:1109.5213, 2011.

The Mathematical Institute, 24-29 St. Giles, Oxford, OX1 3LB, U.K.
E-mail: `joyce@maths.ox.ac.uk`